

Summary of Product Characteristics

1. Name of the medicinal product

Breathezy L

Montelukast Sodium 10 mg and Levocetirizine Dihydrochloride 5 mg Tablets

2. Qualitative and quantitative composition

Each film coated tablet contains

Montelukast sodium Ph.Eur

equivalent to Montelukast 10 mg

Levocetirizine Dihydrochloride 5 mg

For a full list of excipients, see section 6.1.

3. Pharmaceutical form

Film coated tablet

Round shaped, film coated bi layered tablets, one layer with yellow colour and another layer with white to off white color having plain surface on both sides

4. Clinical particulars

4.1 Therapeutic indications

Montelukast is indicated in the treatment of asthma as add-on therapy in those patients with mild to moderate persistent asthma who are inadequately controlled on inhaled corticosteroids and in whom “as-needed” short acting β -agonists provide inadequate clinical control of asthma. In those asthmatic patients in whom Montelukast is indicated in asthma, Montelukast can also provide symptomatic relief of seasonal allergic rhinitis.

Montelukast is also indicated in the prophylaxis of asthma in which the predominant component is exercise-induced bronchoconstriction.

Levocetirizine dihydrochloride indicated in the symptomatic treatment of allergic rhinitis (including persistent allergic rhinitis) and urticaria in adults and children aged 6 years and above.

4.2 Posology and method of administration

Montelukast

Posology

The recommended dose for adults and adolescents 15 years of age and older with asthma, or with asthma and concomitant seasonal allergic rhinitis, is one 10 mg tablet daily to be taken in the evening.

General recommendations

The therapeutic effect of Montelukast on parameters of asthma control occurs within one day. Montelukast may be taken with or without food. Patients should be advised to continue taking Montelukast even if their asthma is under control, as well as during periods of worsening asthma. Montelukast should not be used concomitantly with other products containing the same active ingredient, montelukast.

No dosage adjustment is necessary for the elderly, or for patients with renal insufficiency, or mild to moderate hepatic impairment. There are no data on patients with severe hepatic impairment. The dosage is the same for both male and female patients.

Therapy with Montelukast in relation to other treatments for asthma

Montelukast can be added to a patient's existing treatment regimen.

Inhaled corticosteroids: Treatment with Montelukast can be used as add-on therapy in patients when inhaled corticosteroids plus “as needed” short acting β -agonists provide inadequate clinical control. Montelukast should not be abruptly substituted for inhaled corticosteroids (see section 4.4).

Paediatric population

Do not give Montelukast 10 mg film-coated tablets to children less than 15 years of age. The safety and efficacy of Montelukast 10 mg film-coated tablets in children less than 15 years has not been established.

5 mg chewable tablets are available for paediatric patients 6 to 14 years of age.

4 mg chewable tablets are available for paediatric patients 2 to 5 years of age.

Levocetirizine

Posology

Adults and adolescents 12 years and above:

The daily recommended dose is 5 mg (1 film-coated tablet).

Elderly

Adjustment of the dose is recommended in elderly patients with moderate to severe renal impairment (see Renal impairment below).

Renal impairment

The dosing intervals must be individualised according to renal function (eGFR – estimated Glomerular Filtration Rate). Refer to the following table and adjust the dose as indicated.

Dosing adjustments for patients with impaired renal function:

Group	eGFR (ml/min)	Dosage and frequency
Normal renal function	≥ 90	1 tablet once daily
Mildly decreased renal function	$60 - < 90$	1 tablet once daily
Moderately decreased renal function	$30 - < 60$	1 tablet once every 2 days
Severely decreased renal function	$15 - < 30$ (not requiring dialysis)	1 tablet once every 3 days
End stage renal disease (ESRD)	< 15 (requiring dialysis treatment)	Contra-indicated

In paediatric patients suffering from renal impairment, the dose will have to be adjusted on an individual basis taking into account the renal clearance of the patient and his body weight. There are no specific data for children with renal impairment.

Hepatic impairment

No dose adjustment is needed in patients with solely hepatic impairment. In patients with hepatic impairment and renal impairment, adjustment of the dose is recommended (see Renal impairment above).

Paediatric population

Children aged 6 to 12 years:

The daily recommended dose is 5 mg (1 film-coated tablet).

For children aged 2 to 6 years no adjusted dosage is possible with the film-coated tablet formulation. It is recommended to use a paediatric formulation of levocetirizine.

Method of administration

The film-coated tablet must be taken orally, swallowed whole with liquid and may be taken with or without food. It is recommended to take the daily dose in one single intake.

Duration of use:

Intermittent allergic rhinitis (symptoms experienced for less than four days a week or for less than four weeks a year) has to be treated according to the disease and its history; it can be stopped once the symptoms have disappeared and can be restarted again when symptoms reappear. In case of persistent allergic rhinitis (symptoms experienced for more than four days a week or for more than

four weeks a year), continuous therapy can be proposed to the patient during the period of exposure to allergens.

There is clinical experience with the use of levocetirizine for treatment periods of at least 6 months. In chronic urticaria and chronic allergic rhinitis, there is clinical experience of use of cetirizine (racemate) for up to one year.

4.3 Contraindications

Montelukast

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Levocetirizine

Hypersensitivity to the active substance, to cetirizine, to hydroxyzine, to any other piperazine derivatives or to any of the other excipients listed in section 6.1.

Patients with end stage renal disease with estimated Glomerular Filtration Rate (eGFR) below 15 ml/min (requiring dialysis treatment)

4.4 Special warnings and precautions for use

Montelukast:

Patients should be advised never to use oral montelukast to treat acute asthma attacks and to keep their usual appropriate rescue medication for this purpose readily available. If an acute attack occurs, a short-acting inhaled β -agonist should be used. Patients should seek their doctors' advice as soon as possible if they need more inhalations of short-acting β -agonists than usual.

Montelukast should not be substituted abruptly for inhaled or oral corticosteroids.

There are no data demonstrating that oral corticosteroids can be reduced when montelukast is given concomitantly.

In rare cases, patients on therapy with anti-asthma agents including montelukast may present with systemic eosinophilia, sometimes presenting with clinical features of vasculitis consistent with Churg-Strauss syndrome, a condition which is often treated with systemic corticosteroid therapy. These cases have been sometimes associated with the reduction or withdrawal of oral corticosteroid therapy. Although a causal relationship with leukotriene receptor antagonism has not been established, physicians should be alert to eosinophilia, vasculitic rash, worsening pulmonary symptoms, cardiac complications, and/or neuropathy presenting in their patients. Patients who develop these symptoms should be reassessed and their treatment regimens evaluated.

Treatment with montelukast does not alter the need for patients with aspirin-sensitive asthma to avoid taking aspirin and other non-steroidal anti-inflammatory drugs.

Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Neuropsychiatric events have been reported in adults, adolescents, and children taking Montelukast (see section 4.8). Patients and physicians should be alert for neuropsychiatric events. Patients and/or caregivers should be instructed to notify their physician if these changes occur. Prescribers should carefully evaluate the risks and benefits of continuing treatment with Montelukast if such events occur.

Levocetirizine

Precaution is recommended with concurrent intake of alcohol (see section 4.5).

Caution should be taken in patients with predisposing factors of urinary retention (e.g. spinal cord lesion, prostatic hyperplasia) as levocetirizine may increase the risk of urinary retention.

Caution should be taken in patients with epilepsy and patients at risk of convulsion as levocetirizine may cause seizure aggravation.

Response to allergy skin tests are inhibited by antihistamines and a wash-out period (of 3 days) is required before performing them.

Patients with rare hereditary problems of galactose intolerance, total - lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Pruritus may occur when levocetirizine is stopped even if those symptoms were not present before treatment initiation. The symptoms may resolve spontaneously. In some cases, the symptoms may be intense and may require treatment to be restarted. The symptoms should resolve when the treatment is restarted.

Paediatric population

The use of the film-coated tablet formulation is not recommended in children aged less than 6 years since this formulation does not allow for appropriate dose adaptation. It is recommended to use a paediatric formulation of levocetirizine

4.5 Interaction with other medicinal products and other forms of interaction

Montelukast

Montelukast may be administered with other therapies routinely used in the prophylaxis and chronic treatment of asthma. In drug-interactions studies, the recommended clinical dose of montelukast did not have clinically important effects on the pharmacokinetics of the following medicinal products:

theophylline, prednisone, prednisolone, oral contraceptives (ethinyl estradiol/norethindrone 35/1), terfenadine, digoxin and warfarin.

The area under the plasma concentration curve (AUC) for montelukast was decreased approximately 40% in subjects with co-administration of phenobarbital. Since montelukast is metabolised by CYP 3A4, 2C8, and 2C9, caution should be exercised, particularly in children, when montelukast is co-administered with inducers of CYP 3A4, 2C8, and 2C9, such as phenytoin, phenobarbital and rifampicin.

In vitro studies have shown that montelukast is a potent inhibitor of CYP 2C8. However, data from a clinical drug-drug interaction study involving montelukast and rosiglitazone (a probe substrate representative of medicinal products primarily metabolized by CYP 2C8) demonstrated that montelukast does not inhibit CYP 2C8 *in vivo*. Therefore, montelukast is not anticipated to markedly alter the metabolism of medicinal products metabolised by this enzyme (e.g., paclitaxel, rosiglitazone, and repaglinide).

In vitro studies have shown that montelukast is a substrate of CYP 2C8, and to a less significant extent, of 2C9, and 3A4. In a clinical drug-drug interaction study involving montelukast and gemfibrozil (an inhibitor of both CYP 2C8 and 2C9) gemfibrozil increased the systemic exposure of montelukast by 4.4-fold. No routine dosage adjustment of montelukast is required upon co-administration with gemfibrozil or other potent inhibitors of CYP 2C8, but the physician should be aware of the potential for an increase in adverse reactions.

Based on *in vitro* data, clinically important drug interactions with less potent inhibitors of CYP 2C8 (e.g., trimethoprim) are not anticipated. Co-administration of montelukast with itraconazole, a strong inhibitor of CYP 3A4, resulted in no significant increase in the systemic exposure of montelukast.

Levocetirizine

No interaction studies have been performed with levocetirizine (including no studies with CYP3A4 inducers); studies with the racemate compound cetirizine demonstrated that there were no clinically relevant adverse interactions (with antipyrine, azithromycin, cimetidine, diazepam, erythromycin, glipizide, ketoconazole and pseudoephedrine). A small decrease in the clearance of cetirizine (16%) was observed in a multiple dose study with theophylline (400 mg once a day); while the disposition of theophylline was not altered by concomitant cetirizine administration.

In a multiple dose study of ritonavir (600 mg twice daily) and cetirizine (10 mg daily), the extent of exposure to cetirizine was increased by about 40% while the disposition of ritonavir was slightly altered (-11%) further to concomitant cetirizine administration.

The extent of absorption of levocetirizine is not reduced with food, although the rate of absorption is decreased.

In sensitive patients, the concurrent administration of cetirizine or levocetirizine and alcohol or other CNS depressants may cause additional reductions in alertness and impairment of performance.

4.6 Fertility, pregnancy and lactation

Montelukast

Pregnancy

Animal studies do not indicate harmful effects with respect to effects on pregnancy or embryonal/foetal development.

Available data from published prospective and retrospective cohort studies with montelukast use in pregnant women evaluating major birth defects have not established a drug-associated risk. Available studies have methodologic limitations, including small sample size, in some cases retrospective data collection, and inconsistent comparator groups.

Montelukast may be used during pregnancy only if it is considered to be clearly essential.

Breast-feeding

Studies in rats have shown that montelukast is excreted in milk (see section 5.3). It is unknown whether montelukast/metabolites are excreted in human milk.

Montelukast may be used in breast-feeding only if it is considered to be clearly essential.

Levocetirizine

Pregnancy

There are no or limited amount of data (less than 300 pregnancy outcomes) from the use of levocetirizine in pregnant women. However, for cetirizine, the racemate of levocetirizine, a large amount of data (more than 1000 pregnancy outcomes) on pregnant women indicate no malformative or feto/ neonatal toxicity. Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryo/fetal development, parturition or postnatal development (see section 5.3).

The use of levocetirizine may be considered during pregnancy, if necessary.

Breast-feeding

Cetirizine, the racemate of levocetirizine, has been shown to be excreted in human. Therefore, the excretion of levocetirizine in human milk is likely. Adverse reactions associated with levocetirizine may be observed in breastfed infants. Therefore, caution should be exercised when prescribing levocetirizine to lactating women.

Fertility

For levocetirizine no clinical data are available.

4.7 Effects on ability to drive and use machines

Montelukast

Montelukast has no or negligible influence on the ability to drive and use machines. However, individuals have reported drowsiness or dizziness.

Levocetirizine

recommended dose impairs mental alertness, reactivity or the ability to drive and use machines.

Nevertheless, some patients could experience somnolence, fatigue and asthenia under therapy with levocetirizine. Therefore, patients intending to drive, engage in potentially hazardous activities or operate machinery should take their response to the medicinal product into account.

4.8 Undesirable effects

Montelukast

Montelukast has been evaluated in clinical studies as follows:

- 10 mg film-coated tablets in approximately 4,000 adult and adolescent asthmatic patients 15 years of age and older.
- 10 mg film-coated tablets in approximately 400 adult and adolescent asthmatic patients with seasonal allergic rhinitis 15 years of age and older.
- 5 mg chewable tablets in approximately 1,750 paediatric asthmatic patients 6 to 14 years of age.

The following drug-related adverse reactions in clinical studies were reported commonly ($\geq 1/100$ to $< 1/10$) in asthmatic patients treated with montelukast and at a greater incidence than in patients treated with placebo:

Body System Class	Adult and Adolescent Patients 15 years and older (two 12-week studies; n=795)	Paediatric Patients 6 to 14 years old (one 8-week study; n=201) (two 56-week studies; n=615)
Nervous system disorders	headache	headache
Gastro-intestinal disorders	abdominal pain	

With prolonged treatment in clinical trials with a limited number of patients for up to 2 years for adults, and up to 12 months for paediatric patients 6 to 14 years of age, the safety profile did not change.

Tabulated list of Adverse Reactions

Adverse reactions reported in post-marketing use are listed, by System Organ Class and specific Adverse Reactions, in the table below. Frequency Categories were estimated based on relevant clinical trials.

System Organ Class	Adverse Reactions	Frequency Category*
Infections and infestations	upper respiratory infection [†]	Very Common
Blood and lymphatic system disorders	increased bleeding tendency	Rare
	thrombocytopenia	Very Rare
Immune system disorders	hypersensitivity reactions including anaphylaxis	Uncommon
	hepatic eosinophilic infiltration	Very Rare
Psychiatric disorders	dream abnormalities including nightmares, insomnia, somnambulism, anxiety, agitation including aggressive behaviour or hostility, depression, psychomotor hyperactivity (including irritability, restlessness, tremor [§])	Uncommon
	disturbance in attention, memory impairment, tic	Rare
	hallucinations, disorientation, suicidal thinking and behaviour (suicidality), obsessive-compulsive symptoms, dysphemia	Very Rare
Nervous system disorders	dizziness, drowsiness, paraesthesia/hypoesthesia, seizure	Uncommon
Cardiac disorders	palpitations	Rare
Respiratory, thoracic and mediastinal disorders	epistaxis	Uncommon
	Churg-Strauss Syndrome (CSS) (see section 4.4)	Very Rare

	pulmonary eosinophilia	Very Rare
Gastro-intestinal disorders	diarrhoea [‡] , nausea [‡] , vomiting [‡]	Common
	dry mouth, dyspepsia	Uncommon
Hepatobiliary disorders	elevated levels of serum transaminases (ALT, AST)	Common
	hepatitis (including cholestatic, hepatocellular, and mixed-pattern liver injury).	Very Rare
Skin and subcutaneous tissue disorders	rash [‡]	Common
	bruising, urticaria, pruritus	Uncommon
	angiooedema	Rare
	erythema nodosum, erythema multiforme	Very Rare
Musculoskeletal and connective tissue disorders	arthralgia, myalgia including muscle cramps	Uncommon
Renal and urinary disorders	enuresis in children	Uncommon
General disorders and administration site conditions	pyrexia [‡]	Common
	asthenia/fatigue, malaise, oedema	Uncommon

*Frequency Category: Defined for each Adverse Reaction by the incidence reported in the clinical trials data base: Very Common ($\geq 1/10$), Common ($\geq 1/100$ to $< 1/10$), Uncommon ($\geq 1/1,000$ to $< 1/100$), Rare ($\geq 1/10,000$ to $< 1/1,000$), Very Rare ($< 1/10,000$).

[†]This adverse experience, reported as Very Common in the patients who received montelukast, was also reported as Very Common in the patients who received placebo in clinical trials.

[‡]This adverse experience, reported as Common in the patients who received montelukast, was also reported as Common in the patients who received placebo in clinical trials.

[§] Frequency Category: Rare

Levocetirizine

Clinical studies

Adults and adolescents above 12 years of age

In therapeutic studies in women and men aged 12 to 71 years, 15.1% of the patients in the levocetirizine 5 mg group had at least one adverse drug reaction compared to 11.3% in the placebo group. 91.6 % of these adverse drug reactions were mild to moderate.

In therapeutic trials, the dropout rate due to adverse events was 1.0% (9/935) with levocetirizine 5 mg and 1.8% (14/771) with placebo.

Clinical therapeutic trials with levocetirizine included 935 subjects exposed to the medicinal product at the recommended dose of 5 mg daily. From this pooling, following incidence of adverse drug reactions were reported at rates of 1% or greater (common: $\geq 1/100$ to $< 1/10$) under levocetirizine 5 mg or placebo:

Preferred Term (WHOART)	Placebo (n =771)	Levocetirizine 5 mg (n = 935)
Headache	25 (3.2%)	24 (2.6%)
Somnolence	11 (1.4%)	49 (5.2%)
Mouth dry	12 (1.6%)	24 (2.6%)
Fatigue	9 (1.2%)	23 (2.5%)

Further uncommon incidences of adverse reactions (uncommon $\geq 1/1,000$ to $< 1/100$) like asthenia or abdominal pain were observed.

The incidence of sedating adverse drug reactions such as somnolence, fatigue, and asthenia was altogether more common (8.1%) under levocetirizine 5 mg than under placebo (3.1%).

Paediatric population

In two placebo-controlled studies in paediatric patients aged 6-11 months and aged 1 year to less than 6 years, 159 subjects were exposed to levocetirizine at the dose of 1.25 mg daily for 2 weeks and 1.25 mg twice daily respectively. The following incidence of adverse drug reactions was reported at rates of 1% or greater under levocetirizine or placebo.

System Organ Class and Preferred Term	Placebo (n=83)	Levocetirizine (n=159)
Gastrointestinal disorders		
Diarrhoea	0	3(1.9%)
Vomiting	1(1.2%)	1(0.6%)
Constipation	0	2(1.3%)
Nervous system disorders		
Somnolence	2(2.4%)	3(1.9%)

Psychiatric disorders		
Sleep disorder	0	2(1.3%)

In children aged 6-12 years double blind placebo controlled studies were performed where 243 children were exposed to 5 mg levocetirizine daily for variable periods ranging from less than 1 week to 13 weeks. The following incidence of adverse drug reactions was reported at rates of 1% or greater under levocetirizine or placebo.

Preferred Term	Placebo (n=240)	Levocetirizine 5mg (n=243)
Headache	5(2.1%)	2(0.8%)
Somnolence	1(0.4%)	7(2.9%)

Post-marketing experience

Adverse reactions from post-marketing experience are per System Organ Class and per frequency. The frequency is defined as follows: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$), not known (cannot be estimated from the available data).

- Immune system disorders:

Not known: hypersensitivity including anaphylaxis

- Metabolism and nutrition disorders:

Not known: increased appetite

- Psychiatric disorders:

Not known: aggression, agitation, hallucination, depression, insomnia, suicidal ideation, nightmare

- Nervous system disorders:

Not known: convulsion, paraesthesia, dizziness, syncope, tremor, dysgeusia

- Ear and labyrinth disorders:

Not known: vertigo

- Eyes disorders:

Not known: visual disturbances, blurred vision, oculogyration

- Cardiac disorders:

Not known: palpitations, tachycardia

- Respiratory, thoracic and mediastinal disorders:

Not known: dyspnoea

- Gastrointestinal disorders:

Not known: nausea, vomiting, diarrhoea

- Hepatobiliary disorders:

Not known: hepatitis

- Renal and urinary disorders:

Not known: dysuria, urinary retention

- Skin and subcutaneous tissue disorders:

Not known: angioneurotic oedema, fixed drug eruption, pruritus, rash, urticaria

- Musculoskeletal, connective tissues, and bone disorders:

Not known: myalgia, arthralgia

- General disorders and administration site conditions:

Not known: oedema

- Investigations:

Not known: weight increased, abnormal liver function tests

Description of selected adverse reactions

After levocetirizine discontinuation, pruritus has been reported.

4.9 Overdose

Montelukast

In chronic asthma studies, montelukast has been administered at doses up to 200 mg/day to adult patients for 22 weeks and in short term studies, up to 900 mg/day to patients for approximately one week without clinically important adverse experiences.

There have been reports of acute overdose in post-marketing experience and clinical studies with montelukast. These include reports in adults and children with a dose as high as 1,000 mg (approximately 61 mg/kg in a 42 month old child). The clinical and laboratory findings observed were consistent with the safety profile in adults and paediatric patients. There were no adverse experiences in the majority of overdose reports.

Symptoms of overdose

The most frequently occurring adverse experiences were consistent with the safety profile of montelukast and included abdominal pain, somnolence, thirst, headache, vomiting, and psychomotor hyperactivity.

Management of overdose

No specific information is available on the treatment of overdose with montelukast. It is not known whether montelukast is dialysable by peritoneal- or haemodialysis.

Levocetirizine

Symptoms

Symptoms of overdose may include drowsiness in adults. In children, agitation and restlessness may initially occur, followed by drowsiness.

Management of overdoses

There is no known specific antidote to levocetirizine.

Should overdose occur, symptomatic or supportive treatment is recommended. Gastric lavage may be considered shortly after ingestion of the drug. Levocetirizine is not effectively removed by haemodialysis.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Montelukast

Pharmacotherapeutic group: Leukotriene receptor antagonist

ATC-code: R03D C03

Mechanism of action

The cysteinyl leukotrienes (LTC₄, LTD₄, LTE₄) are potent inflammatory eicosanoids released from various cells including mast cells and eosinophils. These important pro-asthmatic mediators bind to cysteinyl leukotriene (CysLT) receptors. The CysLT type-1 (CysLT₁) receptor is found in the human airway (including airway smooth muscle cells and airway macrophages) and on other pro-inflammatory cells (including eosinophils and certain myeloid stem cells). CysLTs have been correlated with the pathophysiology of asthma and allergic rhinitis. In asthma, leukotriene-mediated effects include bronchoconstriction, mucous secretion, vascular permeability, and eosinophil recruitment. In allergic rhinitis, CysLTs are released from the nasal mucosa after allergen exposure during both early- and late-phase reactions and are associated with symptoms of allergic rhinitis. Intranasal challenge with CysLTs has been shown to increase nasal airway resistance and symptoms of nasal obstruction.

Pharmacodynamic effects

Montelukast is an orally active compound which binds with high affinity and selectivity to the CysLT₁ receptor. In clinical studies, montelukast inhibits bronchoconstriction due to inhaled LTD₄ at doses as low as 5 mg. Bronchodilation was observed within 2 hours of oral administration. The bronchodilation effect caused by a β -agonist was additive to that caused by montelukast. Treatment with montelukast inhibited both early- and late-phase bronchoconstriction due to antigen challenge. Montelukast, compared with placebo, decreased peripheral blood eosinophils in adult and paediatric patients. In a separate study, treatment with montelukast significantly decreased

eosinophils in the airways (as measured in sputum) and in peripheral blood while improving clinical asthma control.

Clinical efficacy and safety

In studies in adults, montelukast, 10 mg once daily, compared with placebo, demonstrated significant improvements in morning FEV₁ (10.4% vs 2.7% change from baseline), AM peak expiratory flow rate (PEFR) (24.5 L/min vs 3.3 L/min change from baseline), and significant decrease in total β -agonist use (-26.1% vs -4.6% change from baseline). Improvement in patient-reported daytime and nighttime asthma symptoms scores was significantly better than placebo.

Studies in adults demonstrated the ability of montelukast to add to the clinical effect of inhaled corticosteroid (% change from baseline for inhaled beclomethasone plus montelukast vs beclomethasone, respectively for FEV₁: 5.43% vs 1.04%; β -agonist use: -8.70% vs 2.64%). Compared with inhaled beclomethasone (200 μ g twice daily with a spacer device), montelukast demonstrated a more rapid initial response, although over the 12-week study, beclomethasone provided a greater average treatment effect (% change from baseline for montelukast vs beclomethasone, respectively for FEV₁: 7.49% vs 13.3%; β -agonist use: -28.28% vs -43.89%). However, compared with beclomethasone, a high percentage of patients treated with montelukast achieved similar clinical responses (e.g., 50% of patients treated with beclomethasone achieved an improvement in FEV₁ of approximately 11% or more over baseline while approximately 42% of patients treated with montelukast achieved the same response).

A clinical study was conducted to evaluate montelukast for the symptomatic treatment of seasonal allergic rhinitis in adult and adolescent asthmatic patients 15 years of age and older with concomitant seasonal allergic rhinitis. In this study, montelukast 10 mg tablets administered once daily demonstrated a statistically significant improvement in the Daily Rhinitis Symptoms score, compared with placebo. The Daily Rhinitis Symptoms score is the average of the Daytime Nasal Symptoms score (mean of nasal congestion, rhinorrhea, sneezing, nasal itching) and the Nighttime Symptoms score (mean of nasal congestion upon awakening, difficulty going to sleep, and nighttime awakenings scores). Global evaluations of allergic rhinitis by patients and physicians were significantly improved, compared with placebo. The evaluation of asthma efficacy was not a primary objective in this study.

In an 8-week study in paediatric patients 6 to 14 years of age, montelukast 5 mg once daily, compared with placebo, significantly improved respiratory function (FEV₁ 8.71% vs 4.16% change from baseline; AM PEFR 27.9 L/min vs 17.8 L/min change from baseline) and decreased “as-needed” β -agonist use (-11.7% vs +8.2% change from baseline).

Significant reduction of exercise-induced bronchoconstriction (EIB) was demonstrated in a 12-week study in adults (maximal fall in FEV₁ 22.33% for montelukast vs 32.40% for placebo; time to recovery to within 5% of baseline FEV₁ 44.22 min vs 60.64 min). This effect was consistent throughout the 12-week study period. Reduction in EIB was also demonstrated in a short term study in paediatric patients (maximal fall in FEV₁ 18.27% vs 26.11%; time to recovery to within 5% of baseline FEV₁ 17.76 min vs 27.98 min). The effect in both studies was demonstrated at the end of the once-daily dosing interval.

In aspirin-sensitive asthmatic patients receiving concomitant inhaled and/or oral corticosteroids, treatment with montelukast, compared with placebo, resulted in significant improvement in asthma control (FEV₁ 8.55% vs -1.74% change from baseline and decrease in total β -agonist use -27.78% vs 2.09% change from baseline).

Levocetirizine:

Pharmacotherapeutic group: antihistamine for systemic use, piperazine derivatives, ATC code: R06A E09.

Mechanism of action

Levocetirizine, the (R) enantiomer of cetirizine, is a potent and selective antagonist of peripheral H₁-receptors.

Binding studies revealed that levocetirizine has high affinity for human H₁-receptors (K_i = 3.2 nmol/l). Levocetirizine has an affinity 2-fold higher than that of cetirizine (K_i = 6.3 nmol/l).

Levocetirizine dissociates from H₁-receptors with a half-life of 115 ± 38 min.

After single administration, levocetirizine shows a receptor occupancy of 90% at 4 hours and 57% at 24 hours.

Pharmacodynamic studies in healthy volunteers demonstrate that, at half the dose, levocetirizine has comparable activity to cetirizine, both in the skin and in the nose.

Pharmacodynamic effects

The pharmacodynamic activity of levocetirizine has been studied in randomised, controlled trials:

In a study comparing the effects of levocetirizine 5 mg, desloratadine 5 mg, and placebo on histamine-induced wheal and flare, levocetirizine treatment resulted in significantly decreased wheal and flare formation which was highest in the first 12 hours and lasted for 24 hours, (p<0.001) compared with placebo and desloratadine.

The onset of action of levocetirizine 5 mg in controlling pollen-induced symptoms has been observed at 1 hour post drug intake in placebo controlled trials in the model of the allergen challenge chamber.

In vitro studies (Boyden chambers and cell layers techniques) show that levocetirizine inhibits eotaxin-induced eosinophil transendothelial migration through both dermal and lung cells. A pharmacodynamic experimental study *in vivo* (skin chamber technique) showed three main inhibitory effects of levocetirizine 5 mg in the first 6 hours of pollen-induced reaction, compared with placebo in 14 adult patients: inhibition of VCAM-1 release, modulation of vascular permeability and a decrease in eosinophil recruitment.

Clinical efficacy and safety

The efficacy and safety of levocetirizine has been demonstrated in several double-blind, placebo controlled, clinical trials performed in adult patients suffering from seasonal allergic rhinitis, perennial allergic rhinitis, or persistent allergic rhinitis. Levocetirizine has been shown to significantly improve symptoms of allergic rhinitis, including nasal obstruction in some studies.

A 6-month clinical study in 551 adult patients (including 276 levocetirizine-treated patients) suffering from persistent allergic rhinitis (symptoms present 4 days a week for at least 4 consecutive weeks) and sensitized to house dust mites and grass pollen demonstrated that levocetirizine 5 mg was clinically and statistically significantly more potent than placebo on the relief from the total symptom score of allergic rhinitis throughout the whole duration of the study, without any tachyphylaxis. During the whole duration of the study, levocetirizine significantly improved the quality of life of the patients.

In a placebo-controlled clinical trial including 166 patients suffering from chronic idiopathic urticaria, 85 patients were treated with placebo and 81 patients with levocetirizine 5 mg once daily over six weeks. Treatment with levocetirizine resulted in significant decrease in pruritus severity over the first week and over the total treatment period as compared to placebo. Levocetirizine also resulted in a larger improvement of health-related quality of life as assessed by the Dermatology Life Quality Index as compared to placebo.

Chronic idiopathic urticaria was studied as a model for urticarial conditions. Since histamine release is a causal factor in urticarial diseases, levocetirizine is expected to be effective in providing symptomatic relief for other urticarial conditions, in addition to chronic idiopathic urticaria.

ECGs did not show relevant effects of levocetirizine on QT interval.

Paediatric population

The paediatric safety and efficacy of levocetirizine tablets has been studied in two placebo controlled clinical trials including patients aged 6 to 12 years and suffering from seasonal and perennial allergic rhinitis, respectively. In both trials, levocetirizine significantly improved symptoms and increased health-related quality of life.

In children below the age of 6 years, clinical safety has been established from several short- or long-term therapeutic studies:

- one clinical trial in which 29 children 2 to 6 years of age with allergic rhinitis were treated with levocetirizine 1.25 mg twice daily for 4 weeks
- one clinical trial in which 114 children 1 to 5 years of age with allergic rhinitis or chronic idiopathic urticaria were treated with levocetirizine 1.25 mg twice daily for 2 weeks
- one clinical trial in which 45 children 6 to 11 months of age with allergic rhinitis or chronic idiopathic urticaria were treated with levocetirizine 1.25 mg once daily for 2 weeks
- one long-term (18 months) clinical trial in 255 levocetirizine - treated atopic subjects aged 12 to 24 months at inclusion.

The safety profile was similar to that seen in the short-term studies conducted in children 1 to 5 years of age.

5.2 Pharmacokinetic properties

Montelukast:

Absorption

Montelukast is rapidly absorbed following oral administration. For the 10 mg film-coated tablet, the mean peak plasma concentration (C_{max}) is achieved 3 hours (T_{max}) after administration in adults in the fasted state. The mean oral bioavailability is 64%. The oral bioavailability and C_{max} are not influenced by a standard meal. Safety and efficacy were demonstrated in clinical trials where the 10 mg film-coated tablet was administered without regard to the timing of food ingestion.

For the 5 mg chewable tablet, the C_{max} is achieved in 2 hours after administration in adults in the fasted state. The mean oral bioavailability is 73% and is decreased to 63% by a standard meal.

Distribution

Montelukast is more than 99% bound to plasma proteins. The steady-state volume of distribution of montelukast averages 8-11 litres. Studies in rats with radiolabelled montelukast indicate minimal distribution across the blood-brain barrier. In addition, concentrations of radiolabelled material at 24 hours' post-dose were minimal in all other tissues.

Biotransformation

Montelukast is extensively metabolised. In studies with therapeutic doses, plasma concentrations of metabolites of montelukast are undetectable at steady state in adults and children.

Cytochrome P450 2C8 is the major enzyme in the metabolism of montelukast. Additionally, CYP 3A4 and 2C9 may have a minor contribution, although itraconazole, an inhibitor of CYP 3A4, was

shown not to change pharmacokinetic variables of montelukast in healthy subjects that received 10 mg montelukast daily. Based on *in vitro* results in human liver microsomes, therapeutic plasma concentrations of montelukast do not inhibit cytochromes P450 3A4, 2C9, 1A2, 2A6, 2C19, or 2D6. The contribution of metabolites to the therapeutic effect of montelukast is minimal.

Elimination

The plasma clearance of montelukast averages 45 ml/min in healthy adults. Following an oral dose of radiolabelled montelukast, 86% of the radioactivity was recovered in 5-day faecal collections and <0.2% was recovered in urine. Coupled with estimates of montelukast oral bioavailability, this indicates that montelukast and its metabolites are excreted almost exclusively via the bile.

Characteristics in Patients

No dosage adjustment is necessary for the elderly or mild to moderate hepatic insufficiency. Studies in patients with renal impairment have not been undertaken. Because montelukast and its metabolites are eliminated by the biliary route, no dose adjustment is anticipated to be necessary in patients with renal impairment. There are no data on the pharmacokinetics of montelukast in patients with severe hepatic insufficiency (Child-Pugh score >9).

With high doses of montelukast (20- and 60-fold the recommended adult dose), decrease in plasma theophylline concentration was observed. This effect was not seen at the recommended dose of 10 mg once daily.

Levocetirizine:

The pharmacokinetics of levocetirizine are linear with dose- and time-independent with low inter-subject variability. The pharmacokinetic profile is the same when given as the single enantiomer or when given as cetirizine. No chiral inversion occurs during the process of absorption and elimination.

Absorption

Levocetirizine is rapidly and extensively absorbed following oral administration. In adults, peak plasma concentrations are achieved 0.9 h after dosing. Steady state is achieved after two days. Peak concentrations are typically 270 ng/ml and 308 ng/ml following a single and a repeated 5 mg o.d. dose, respectively. The extent of absorption is dose-independent and is not altered by food, but the peak concentration is reduced and delayed.

Distribution

No tissue distribution data are available in humans, neither concerning the passage of levocetirizine through the blood-brain-barrier. In rats and dogs, the highest tissue levels are found in liver and kidneys, the lowest in the CNS compartment.

In humans, levocetirizine is 90% bound to plasma proteins. The distribution of levocetirizine is restrictive, as the volume of distribution is 0.4 l/kg.

Biotransformation

The extent of metabolism of levocetirizine in humans is less than 14% of the dose and therefore differences resulting from genetic polymorphism or concomitant intake of enzyme inhibitors are expected to be negligible. Metabolic pathways include aromatic oxidation, N- and O- dealkylation and taurine conjugation. Dealkylation pathways are primarily mediated by CYP 3A4 while aromatic oxidation involved multiple and/or unidentified CYP isoforms. Levocetirizine had no effect on the activities of CYP isoenzymes 1A2, 2C9, 2C19, 2D6, 2E1 and 3A4 at concentrations well above peak concentrations achieved following a 5 mg oral dose.

Due to its low metabolism and absence of metabolic inhibition potential, the interaction of levocetirizine with other substances, or vice-versa, is unlikely.

Elimination

The plasma half-life in adults is 7.9 ± 1.9 hours. The half-life is shorter in small children.

The mean apparent total body clearance in adults is 0.63 ml/min/kg. The major route of excretion of levocetirizine and metabolites is via urine, accounting for a mean of 85.4% of the dose. Excretion via faeces accounts for only 12.9% of the dose. Levocetirizine is excreted both by glomerular filtration and active tubular secretion.

Special population

Renal impairment

The apparent body clearance of levocetirizine is correlated to the creatinine clearance. It is therefore recommended to adjust the dosing intervals of levocetirizine in patients with moderate and severe renal impairment (see section 4.2). In anuric end stage renal disease subjects, the total body clearance is decreased by approximately 80% when compared to normal subjects. The amount of levocetirizine removed during a standard 4-hour hemodialysis procedure was < 10%.

Paediatric population

Data from a paediatric pharmacokinetic study with oral administration of a single dose of 5 mg levocetirizine in 14 children age 6 to 11 years with body weight ranging between 20 and 40 kg show that C_{max} and AUC values are about 2-fold greater than that reported in healthy adult subjects in a cross-study comparison. The mean C_{max} was 450 ng/ml, occurring at a mean time of 1.2 hours, weight-normalized, total body clearance was 30% greater, and the elimination half-life 24% shorter in this paediatric population than in adults. Dedicated pharmacokinetic studies have not been conducted in paediatric patients younger than 6 years of age. A retrospective population pharmacokinetic analysis was conducted in 323 subjects (181 children 1 to 5 years of age, 18

children 6 to 11 years of age, and 124 adults 18 to 55 years of age) who received single or multiple doses of levocetirizine ranging from 1.25 mg to 30 mg. Data generated from this analysis indicated that administration of 1.25 mg once daily to children 6 months to 5 years of age is expected to result in plasma concentrations similar to those of adults receiving 5 mg once daily.

Elderly

Limited pharmacokinetic data are available in elderly subjects. Following once daily repeat oral administration of 30 mg levocetirizine for 6 days in 9 elderly subjects (65–74 years of age), the total body clearance was approximately 33% lower compared to that in younger adults. The disposition of racemic cetirizine has been shown to be dependent on renal function rather than on age. This finding would also be applicable for levocetirizine, as levocetirizine and cetirizine are both predominantly excreted in urine. Therefore, the levocetirizine dose should be adjusted in accordance with renal function in elderly patients.

Gender

Pharmacokinetic results for 77 patients (40 men, 37 women) were evaluated for potential effect of gender. The half-life was slightly shorter in women (7.08 ± 1.72 hr) than in men (8.62 ± 1.84 hr); however, the body weight-adjusted oral clearance in women (0.67 ± 0.16 ml/min/kg) appears to be comparable to that in men (0.59 ± 0.12 ml/min/kg). The same daily doses and dosing intervals are applicable for men and women with normal renal function.

Race

The effect of race on levocetirizine has not been studied. As levocetirizine is primarily renally excreted, and there are no important racial differences in creatinine clearance, pharmacokinetic characteristics of levocetirizine are not expected to be different across races. No race-related differences in the kinetics of racemic cetirizine have been observed.

Hepatic impairment

The pharmacokinetics of levocetirizine in hepatically impaired subjects have not been tested. Patients with chronic liver diseases (hepatocellular, cholestatic, and biliary cirrhosis) given 10 or 20 mg of the racemic compound cetirizine as a single dose had a 50% increase in half life along with a 40% decrease in clearance compared to healthy subjects.

Pharmacokinetic / pharmacodynamic relationship

The action on histamine-induced skin reactions is out of phase with the plasma concentrations.

5.3 Preclinical safety data

Montelukast:

In animal toxicity studies, minor serum biochemical alterations in ALT, glucose, phosphorus and triglycerides were observed which were transient in nature. The signs of toxicity in animals were increased excretion of saliva, gastrointestinal symptoms, loose stools and ion imbalance. These occurred at dosages which provided >17-fold the systemic exposure seen at the clinical dosage. In monkeys, the adverse effects appeared at doses from 150 mg/kg/day (>232-fold the systemic exposure seen at the clinical dose). In animal studies, montelukast did not affect fertility or reproductive performance at systemic exposure exceeding the clinical systemic exposure by greater than 24-fold. A slight decrease in pup body weight was noted in the female fertility study in rats at 200 mg/kg/day (>69-fold the clinical systemic exposure). In studies in rabbits, a higher incidence of incomplete ossification, compared with concurrent control animals, was seen at systemic exposure >24-fold the clinical systemic exposure seen at the clinical dose. No abnormalities were seen in rats. Montelukast has been shown to cross the placental barrier and is excreted in breast milk of animals. No deaths occurred following a single oral administration of montelukast sodium at doses up to 5,000 mg/kg in mice and rats (15,000 mg/m² and 30,000 mg/m² in mice and rats, respectively), the maximum dose tested. This dose is equivalent to 25,000 times the recommended daily adult human dose (based on an adult patient weight of 50 kg).

Montelukast was determined not to be phototoxic in mice for UVA, UVB or visible light spectra at doses up to 500 mg/kg/day (approximately >200-fold based on systemic exposure).

Montelukast was neither mutagenic in *in vitro* and *in vivo* tests nor tumorigenic in rodent species.

Levocetirizine

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction and development.

6. Pharmaceutical particulars

6.1 List of excipients

Core : Lactose monohydrate, Microcrystalline cellulose pH 112, Croscarmellose sodium, Hydroxy propyl cellulose, Magnesium stearate,, Yellow oxide of iron, Colloidal silicon dioxide,
Coating: Instacoat aqua IC-A-329, Purified water

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

2 years

6.4 Special precautions for storage

Do not store above 30°C Protect from moisture. Keep out of reach of children.

6.5 Nature and contents of container

Ten tablets are packed in Aluminium base foil on one side and Aluminium lidding foil another side in the form of Aluminium-Aluminium blister pack. Such three blisters are further packed in pre-printed on carton along with leaflet..

6.6 Special precautions for disposal and other handling

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. Marketing authorisation holder

MSN LABORATORIES PRIVATE LIMITED

“MSN HOUSE” Plot No.C24,

Industrial Estate, Sanathnagar,

Hyderabad 500 018, Telangana, India

Manufactured by

MSN LABORATORIES PRIVATE LIMITED

Plot No: 42, Anrich Industrial Estate,

IDA, Bollaram, Sangareddy Dist,

Telangana,India

8. Marketing authorisation number(s)

05848/07177/NMR/2019

9. Date of first authorisation/renewal of the authorisation

07-04-2021

10. Date of revision of the text

July 2023