

## **SUMMARY OF PRODUCT CHARACTERSTICS**

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE FINISHED PHARMACEUTICAL PRODUCT

**Product Name** : UZET (Azithromycin Tablets USP 500mg)  
**Strength** : 500 mg  
**Pharmaceutical Form** : Tablet

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

#### Label Claim:

Each film-coated tablet contains:  
Azithromycin Dihydrate USP eq. to.  
Anhydrous Azithromycin 500 mg  
Excipients .....q.s.  
Colour: Titanium Dioxide.

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Tablet

White, oblong shaped, film coated, biconvex tablets product plain on both the sides.

### 4. CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

Azithromycin tablets can be applied for the treatment of the following infections, when caused by microorganisms sensitive to azithromycin:

- acute bacterial sinusitis (adequately diagnosed)
- acute bacterial otitis media (adequately diagnosed)
- pharyngitis, tonsillitis
- acute exacerbation of chronic bronchitis (adequately diagnosed)
- mild to moderately severe community acquired pneumonia
- skin and soft tissue infections
- uncomplicated *Chlamydia trachomatis* urethritis and cervicitis

Considerations should be given to official guidance on the appropriate use of antibacterial agents.

#### 4.2 Posology and method of administration

##### Posology

*Adults*: In uncomplicated *Chlamydia trachomatis* urethritis and cervicitis the dose is 1000 mg as a single oral dose.

For all other indications the dose is 1500 mg, to be administered as 500 mg per day for three consecutive days. As an alternative the same total dose (1500 mg) can also be administered over a period of five days with 500 mg on the first day and 250 mg on the second to the fifth day.

#### *Elderly people*

The same dose as in adult patients is used for older people. Since elderly people can be patients with ongoing proarrhythmic conditions a particular caution is recommended due to the risk of developing cardiac arrhythmia and torsades de pointes.

#### *Paediatric population*

Azithromycin tablets should only be administered to children weighing more than 45 kg when normal adult dose should be used. For children under 45 kg other pharmaceutical forms of azithromycine, e.g. suspensions, may be used.

*In patients with renal impairment:* No dose adjustment is necessary in patients with mild to moderate renal impairment (GFR 10-80 ml/min).

*In patients with hepatic impairment:* A dose adjustment is not necessary for patients with mild to moderately impaired liver function.

#### Method of administration

Azithromycin Tablets should be given as a single daily dose. The tablets may be taken with food.

### **4.3 Contraindications**

Hypersensitivity to the active substance, erythromycin, any macrolide, ketolide antibiotic, soya lecithin or to any of the excipients listed.

### **4.4 Special warnings and precautions for use**

#### Hypersensitivity

As with erythromycin and other macrolides, rare serious allergic reactions including angioneurotic oedema and anaphylaxis (rarely fatal), dermatologic reactions including acute generalised exanthematous pustulosis (AGEP), Stevens Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) (rarely fatal) and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported. Some of these reactions with azithromycin have resulted in recurrent symptoms and required a longer period of observation and treatment.

If an allergic reaction occurs, the medicinal product should be discontinued and appropriate therapy should be instituted. Physicians should be aware that reappearance of the allergic symptoms may occur when symptomatic therapy is discontinued.

Since liver is the principal route of elimination for azithromycin, the use of azithromycin should be undertaken with caution in patients with significant hepatic disease. Cases of fulminant hepatitis potentially leading to life-threatening liver failure have been reported with azithromycin. Some patients may have had pre-existing hepatic disease or may have been taking other hepatotoxic medicinal products.

In case of signs and symptoms of liver dysfunction, such as rapid developing asthenia associated with jaundice, dark urine, bleeding tendency or hepatic encephalopathy, liver

function tests / investigations should be performed immediately. Azithromycin administration should be stopped if liver dysfunction has emerged.

In patients receiving ergotamine derivatives, ergotism has been precipitated by coadministration of some macrolide antibiotics. There are no data concerning the possibility of an interaction between ergotamine derivatives and azithromycin. However, because of the theoretical possibility of ergotism, azithromycin and ergot derivatives should not be co-administered.

Because of existing cross-resistance with erythromycin-resistant gram-positive strains and most strains of methicillin resistant staphylococci, use of azithromycin is not recommended.

Local epidemiology and susceptibility patterns should be taken into consideration.

### Cardiovascular events

Prolonged cardiac repolarisation and QT interval, imparting a risk of developing cardiac arrhythmia and torsades de pointes, have been seen in treatment with other macrolides including azithromycin (see section 4.8). Therefore as the following situations may lead to an increased risk for ventricular arrhythmias (including torsade de pointes) which can lead to cardiac arrest, azithromycin should be used with caution in patients with ongoing proarrhythmic conditions (especially women and elderly patients) such as patients:

- With congenital or documented QT prolongation.
- Currently receiving treatment with other active substances known to prolong QT interval such as antiarrhythmics of class IA (quinidine and procainamide) and class III (dofetilide, amiodarone and sotalol), cisapride and terfenadine; antipsychotic agents such as pimozide; antidepressants such as citalopram; and fluoroquinolones such as moxifloxacin and levofloxacin
- With electrolyte disturbance, particularly in cases of hypokalaemia and hypomagnesaemia
- With clinically relevant bradycardia, cardiac arrhythmia or severe cardiac insufficiency.

Epidemiological studies investigating the risk of adverse cardiovascular outcomes with macrolides have shown variable results. Some observational studies have identified a rare short term risk of arrhythmia, myocardial infarction and cardiovascular mortality associated with macrolides including azithromycin. Consideration of these findings should be balanced with treatment benefits when prescribing azithromycin.

*Clostridium difficile* associated diarrhoea (CDAD) has been reported with the use of nearly all antibacterial agents, including azithromycin, and may range in severity from mild diarrhoea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

*C. difficile* produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhoea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antimicrobial agents. In case of CDAD anti-peristaltics are contraindicated.

Exacerbations of the symptoms of myasthenia gravis and new onset of myasthenia syndrome have been reported in patients receiving azithromycin therapy.

Safety and efficacy for the prevention or treatment of Mycobacterium avium complex in children have not been established.

#### Long-term use

There is no experience on safety and effectiveness of long-term use of azithromycin in indications mentioned before. At fast recurrent infections, treatment with other antibiotics should be considered.

#### **The following should be considered before prescribing azithromycin:**

Azithromycin film-coated tablets are not suitable for treatment of severe infections where a high concentration of the antibiotic in the blood is rapidly needed.

Azithromycin is not the first choice for the empiric treatment of infections in areas where the prevalence of resistant isolates is 10% or more.

In areas with a high incidence of erythromycin A resistance, it is especially important to take into consideration the evolution of the pattern of susceptibility to azithromycin and other antibiotics.

As for other macrolides, high resistance rates of *Streptococcus pneumoniae* (> 30 %) have been reported for azithromycin in some European countries. This should be taken into account when treating infections caused by *Streptococcus pneumoniae*.

#### Pharyngitis/ tonsillitis

Azithromycin is not the substance of first choice for the treatment of pharyngitis and tonsillitis caused by *Streptococcus pyogenes*. For this and for the prophylaxis of acute rheumatic fever penicillin is the treatment of first choice.

#### Sinusitis

Often, azithromycin is not the substance of first choice for the treatment of sinusitis.

#### Acute otitis media

Often, azithromycin is not the substance of first choice for the treatment of acute otitis media.

#### Skin and soft tissue infections

The main causative agent of soft tissue infections, *Staphylococcus aureus*, is frequently resistant to azithromycin. Therefore, susceptibility testing is considered a precondition for treatment of soft tissue infections with azithromycin.

#### Infected burn wounds

Azithromycin is not indicated for the treatment of infected burn wounds.

#### Sexually transmitted disease

In case of sexually transmitted diseases a concomitant infection by *T. palladium* should be excluded.

### Neurological or psychiatric diseases

Azithromycin should be used with caution in patients with neurological or psychiatric disorders.

As with any antibiotic preparation, observation for signs of superinfection with non-susceptible organisms, including fungi is recommended.

In patients with severe renal impairment (GFR < 10 ml/min) a 33% increase in systemic exposure to azithromycin was observed.

Azithromycin Tablets contains soya lecithin which might be a source of soya protein and should therefore not be taken in patients allergic to soya or peanut due to the risk of hypersensitivity reactions.

Azithromycin Tablets contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

### **4.5 Interaction with other medicinal products and other forms of interaction** **Effects of other medicinal products on azithromycin:**

#### *Antacids*

In a pharmacokinetic study investigating the effects of simultaneous administration of antacids and azithromycin, no effect on overall bioavailability was seen, although the peak serum concentrations were reduced by approximately 24%. In patients receiving both azithromycin and antacids, the medicinal products should not be taken simultaneously. Azithromycin must be taken at least 1 hour before or 2 hours after the antacids.

Co-administration of azithromycin prolonged-release granules for oral suspension with a single 20 ml dose of co-magaldrox (aluminium hydroxide and magnesium hydroxide) did not affect the rate and extent of azithromycin absorption.

Co-administration of a 600 mg single dose of azithromycin and 400 mg efavirenz daily for 7 days did not result in any clinically significant pharmacokinetic interactions.

#### *Fluconazole*

Co-administration of a single dose of 1200 mg azithromycin did not alter the pharmacokinetics of a single dose of 800 mg fluconazole. Total exposure and half-life of azithromycin were unchanged by the coadministration of fluconazole, however, a clinically insignificant decrease in  $C_{max}$  (18%) of azithromycin was observed.

#### *Nelfinavir*

Co-administration of azithromycin (1200 mg) and nelfinavir at steady state (750 mg three times daily) resulted in increased azithromycin concentrations. No clinically significant adverse effects were observed and no dose adjustment is required.

#### *Rifabutin*

Coadministration of azithromycin and rifabutin did not affect the serum concentrations of either medicinal product.

Neutropenia was observed in subjects receiving concomitant treatment of azithromycin and rifabutin. Although neutropenia has been associated with the use of rifabutin, a causal relationship to combination with azithromycin has not been established.

#### *Terfenadine*

Pharmacokinetic studies have reported no evidence of an interaction between azithromycin and terfenadine. There have been rare cases reported where the possibility of such an interaction could not be entirely excluded; however there was no specific evidence that such an interaction had occurred.

#### *Cimetidine*

In a pharmacokinetic study investigating the effects of a single dose of cimetidine, given 2 hours before azithromycin, on the pharmacokinetics of azithromycin, no alteration of azithromycin pharmacokinetics was seen.

### **Effect of azithromycin on other medicinal products:**

#### *Ergotamine derivatives*

Due to the theoretical possibility of ergotism, the concurrent use of azithromycin with ergot derivatives is not recommended.

#### *Digoxin and colchicine (P-gp substrates)*

Concomitant administration of macrolide antibiotics, including azithromycin, with P-glycoprotein substrates such as digoxin and colchicine, has been reported to result in increased serum levels of the P-glycoprotein substrate. Therefore, if azithromycin and P-gp substrates such as digoxin are administered concomitantly, the possibility of elevated serum concentrations of the substrate should be considered.

#### *Coumarin-Type Oral Anticoagulants*

In a pharmacokinetic interaction study, azithromycin did not alter the anticoagulant effect of a single 15-mg dose of warfarin administered to healthy volunteers. There have been reports received in the post-marketing period of potentiated anticoagulation subsequent to co-administration of azithromycin and coumarin-type oral anticoagulants. Although a causal relationship has not been established, consideration should be given to the frequency of monitoring prothrombin time when azithromycin is used in patients receiving coumarin-type oral anticoagulants.

#### *Cyclosporin*

In a pharmacokinetic study with healthy volunteers that were administered a 500 mg/day oral dose of azithromycin for 3 days and were then administered a single 10 mg/kg oral dose of cyclosporin, the resulting cyclosporin  $C_{max}$  and  $AUC_{0-5}$  were found to be significantly elevated. Consequently, caution should be exercised before considering concurrent administration of these drugs. If coadministration of these drugs is necessary, cyclosporin levels should be monitored and the dose adjusted accordingly.

#### *Theophylline*

There is no evidence of a clinically significant pharmacokinetic interaction when azithromycin and theophylline are co-administered to healthy volunteers. As interactions of

other macrolides with theophylline have been reported, alertness to signs that indicate a rise in theophylline levels is advised.

#### *Trimethoprim/sulfamethoxazole*

Coadministration of trimethoprim/sulfamethoxazole DS (160 mg/800 mg) for 7 days with azithromycin 1200 mg on Day 7 had no significant effect on peak concentrations total exposure or urinary excretion of either trimethoprim or sulfamethoxazole. Azithromycin serum concentrations were similar to those seen in other studies.

#### *Zidovudine*

Single 1000 mg doses and multiple 1200 mg or 600 mg doses of azithromycin had little effect on the plasma pharmacokinetics or urinary excretion of zidovudine or its glucuronide metabolite. However, administration of azithromycin increased the concentrations of phosphorylated zidovudine, the clinically active metabolite, in peripheral blood mononuclear cells. The clinical significance of this finding is unclear, but it may be of benefit to patients.

Azithromycin does not interact significantly with the hepatic cytochrome P450 system. It is not believed to undergo the pharmacokinetic drug interactions as seen with erythromycin and other macrolides. Hepatic cytochrome P450 induction or inactivation via cytochrome-metabolite complex does not occur with azithromycin.

#### *Astemizole, alfentanil*

There are no known data on interactions with astemizole or alfentanil. Caution is advised in the co-administration of these medicines with azithromycin because of the known enhancing effect of these medicines when used concurrently with the macrolid antibiotic erythromycin.

#### *Atorvastatin*

Coadministration of atorvastatin (10 mg daily) and azithromycin (500 mg daily) did not alter the plasma concentrations of atorvastatin (based on a HMG CoA-reductase inhibition assay).

However, post-marketing cases of rhabdomyolysis in patients receiving azithromycin with statins have been reported.

#### *Carbamazepine*

In a pharmacokinetic interaction study in healthy volunteers, no significant effect was observed on the plasma levels of carbamazepine or its active metabolite in patients receiving concomitant azithromycin.

#### *Cisapride*

Cisapride is metabolized in the liver by the enzyme CYP 3A4. Because macrolides inhibit this enzyme, concomitant administration of cisapride may cause the increase of QT interval prolongation, ventricular arrhythmias and torsades de pointes.

#### *Cetirizine*

In healthy volunteers, coadministration of a 5-day regimen of azithromycin with cetirizine 20 mg at steady-state resulted in no pharmacokinetic interaction and no significant changes in the QT interval.



### *Didanosins (Dideoxyinosine)*

Coadministration of 1200 mg/day azithromycin with 400 mg/day didanosine in 6 HIV-positive subjects did not appear to affect the steady-state pharmacokinetics of didanosine as compared with placebo.

### *Efavirenz*

Coadministration of a 600 mg single dose of azithromycin and 400 mg efavirenz daily for 7 days did not result in any clinically significant pharmacokinetic interactions.

### *Indinavir*

Coadministration of a single dose of 1200 mg azithromycin had no statistically significant effect on the pharmacokinetics of indinavir administered as 800 mg three times daily for 5 days.

### *Methylprednisolone*

In a pharmacokinetic interaction study in healthy volunteers, azithromycin had no significant effect on the pharmacokinetics of methylprednisolone.

### *Midazolam*

In healthy volunteers, coadministration of azithromycin 500 mg/day for 3 days did not cause clinically significant changes in the pharmacokinetics and pharmacodynamics of a single 15 mg dose of midazolam.

### *Sildenafil*

In normal healthy male volunteers, there was no evidence of an effect of azithromycin (500 mg daily for 3 days) on the AUC and  $C_{max}$  of sildenafil or its major circulating metabolite.

### *Triazolam*

In 14 healthy volunteers, coadministration of azithromycin 500 mg on Day 1 and 250 mg on Day 2 with 0.125 mg triazolam on Day 2 had no significant effect on any of the pharmacokinetic variables for triazolam compared to triazolam and placebo.

### Other antibiotics

On a possible co-resistance between macrolide antibiotics and azithromycin (e.g. erythromycin) as well as lincomycin and clindamycin is to look at. Concomitant use of several medicinal products from the same group of substances is not recommended.

## **4.6 Fertility, pregnancy and lactation**

### *Pregnancy*

There are no adequate data from the use of azithromycin in pregnant women. In reproduction toxicity studies in animals azithromycin was shown to pass the placenta, but no teratogenic effects were observed. The safety of azithromycin has not been confirmed with regard to the use of the active substance during pregnancy. Therefore azithromycin should only be used during pregnancy if the benefit outweighs the risk.

### *Breast-feeding*

Azithromycin has been reported to be secreted into human breast milk, but there are no adequate and well-controlled clinical studies in nursing women that have characterized the pharmacokinetics of azithromycin excretion into human breast milk.

Because it is not known whether azithromycin may have adverse effects on the breast-fed infant, nursing should be discontinued during treatment with azithromycin. Among other things diarrhoea, fungus infection of the mucous membrane as well as sensitisation is possible in the nursed infant. It is recommended to discard the milk during treatment and up until 2 days after discontinuation of treatment. Nursing may be resumed thereafter.

### *Fertility*

In fertility studies conducted in rat, reduced pregnancy rates were noted following administration of azithromycin. The relevance of this finding to humans is unknown.

#### **4.7 Effects on ability to drive and use machines**

There is no evidence to suggest that azithromycin may have an effect: on a patient's ability to drive or operate machinery. However, certain adverse reactions that have been reported with azithromycin may affect some patients' ability to drive or operate machinery. Visual impairment and vision blurred may have an effect on a patient's ability to drive or operate machinery.

#### **4.8 Undesirable effects**

The table below lists the adverse reactions identified through clinical experience and post-marketing surveillance by system organ class and frequency. Adverse reactions identified from post-marketing experience are included in italics. The frequency grouping is defined using the following convention: Very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Approximately 13% of patients in clinical trials reported adverse events, wherein gastrointestinal disorders were the most.

#### **Adverse reactions possibly or probably related to azithromycin based on clinical trial experience and post-marketing surveillance:**

<b>System Organ Class</b>	<b>Frequency</b>	<b>Adverse reaction</b>
Infections and infestations	Uncommon	Candidiasis Vaginal infection Pneumonia Fungal infection Bacterial infection Pharyngitis Gastroenteritis Respiratory disorder Rhinitis Oral candidiasis
	Not known	<i>Pseudomembranous colitis</i>
Blood and lymphatic system disorders	Uncommon	Leukopenia Neutropenia

		Eosinophilia
	Not known	Thrombocytopenia Haemolytic anaemia
Immune system disorders	Uncommon	Angioedema Hypersensitivity
	Not known	Anaphylactic reaction
Metabolism and nutrition disorders	Uncommon	Anorexia
Psychiatric disorders	Uncommon	Nervousness Insomnia
	Rare	Agitation Depersonalisation
	Not known	<i>Aggression</i> <i>Anxiety</i> Delirium Hallucination
Nervous system disorders	Common	Headache
	Uncommon	Dizziness Somnolence Dysgeusia Paraesthesia
	Not known	<i>Syncope, convulsion</i> Hypoaesthesia <i>Psychomotor hyperactivity</i> <i>Anosmia</i> <i>Ageusia</i> <i>Parosmia</i> <i>Myasthenia gravis</i>
Eye disorders	Uncommon	Visual impairment
	Not known	Blurred vision
Ear and labyrinth disorders	Uncommon	Ear disorder Vertigo
	Not known	Hearing impairment including deafness and/or tinnitus
Cardiac disorders	Uncommon	Palpitations
	Not known	<i>Torsades de pointes</i> <i>Arrhythmia including ventricular tachycardia</i> <i>electrocardiogram QT prolonged</i>
Vascular disorders	Uncommon	Hot flush
	Not known	<i>Hypotension</i>
Respiratory, thoracic and mediastinal disorders	Uncommon	Dyspnoea Epistaxis
Gastrointestinal disorders	Very common	Diarrhoea

	Common	Vomiting Abdominal pain Nausea
	Uncommon	Constipation Flatulence Dyspepsia Gastritis Dysphagia Abdominal distension Dry mouth Eructation Mouth ulceration Salivary hypersecretion
	Not known	<i>Pancreatitis</i> <i>Tongue discolouration</i>
Hepatobiliary disorders	Uncommon	Hepatitis
	Rare	Hepatic function abnormal Jaundice cholestatic
	Not known	<i>Hepatic failure</i> (which has rarely resulted in death)* <i>Hepatitis fulminant</i> <i>Hepatic necrosis</i>
Skin and subcutaneous tissue disorders	Uncommon	Rash Pruritus Urticaria Dermatitis Dry skin Hyperhidrosis
	Rare	Photosensitivity reaction Acute generalised exanthematous pustulosis (AGEP)
	Not known	Steven-Johnson syndrome <i>Toxic epidermal necrolysis</i> <i>Erythema multiforme</i>
Musculoskeletal and connective tissue disorders	Uncommon	Osteoarthritis Myalgia Back pain Neck pain
	Not known	Arthralgia
Renal and urinary disorders	Uncommon	Dysuria Renal pain
	Not known	Renal failure acute Nephritis interstitial
Reproductive system and breast disorders	Uncommon	Metrorrhagia Testicular disorder
General disorders and	Uncommon	Oedema

administration site conditions		Asthenia Malaise Fatigue Face oedema Chest pain Pyrexia Pain Peripheral oedema
Investigations	Common	Lymphocyte count decreased Eosinophil count increased Blood bicarbonate decreased Basophils increased Monocytes increased Neutrophils increased
	Uncommon	Aspartate aminotransferase increased Alanine aminotransferase increased Blood bilirubine increased Blood urea increased Blood creatinine increased Blood potassium abnormal Blood alkaline phosphatase increased Chloride increased Glucose increased Platelets increased Hematocrit decreased Bicarbonate increased Abnormal sodium
Injury and poisoning	Uncommon	Post procedural complication

\* which has rarely resulted in death

**Adverse reactions possibly or probably related to Mycobacterium Avium Complex prophylaxis and treatment based on clinical trial experience and post-marketing surveillance. These adverse reactions differ from those reported with immediate release or the prolonged release formulations, either in kind or in frequency:**

System Organ Class	Frequency	Adverse reaction
Metabolism and nutrition disorders	Common	Anorexia
Nervous system disorders	Common	Dizziness Headache Paraesthesia Dysgeusia
	Uncommon	Hypoaesthesia
Eye disorders	Common	Visual impairment
Ear and labyrinth disorders	Common	Deafness
	Uncommon	Hearing impaired

		Tinnitus
Cardiac disorders	Uncommon	Palpitations
Gastrointestinal disorders	Very common	Diarrhoea Abdominal pain Nausea Flatulence Abdominal discomfort Loose stools
Hepatobiliary disorders	Uncommon	Hepatitis
Skin and subcutaneous tissue disorders	Common	Rash Pruritus
	Uncommon	Steven-Johnson syndrome Photosensitivity reaction
Musculoskeletal and connective tissue disorders	Common	Arthralgia
General disorders and administration site conditions	Common	Fatigue
	Uncommon	Asthenia Malaise

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

#### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via EFDA yellow Card Scheme, online at <https://primaryreporting.who-umc.org/ET> or toll free call 8482 to Ethiopian food and drug authority (EFDA).

#### **4.9 Overdose**

Adverse events experienced in higher than recommended doses were similar to those seen at normal doses. In the event of overdose general symptomatic and general supportive measures are indicated as required.

### **5. Pharmacological properties**

#### **5.1 Pharmacodynamic properties**

##### **General properties**

Pharmacotherapeutic group: antibacterials for systemic use; macrolids; azithromycin, ATC code: J01FA10

##### Mode of action:

Azithromycin is an azalide, a sub-class of the macrolid antibiotics. By binding to the 50S-ribosomal sub-unit, azithromycin avoids the translocation of peptide chains from one side of the ribosome to the other. As a consequence of this, RNA-dependent protein synthesis in sensitive organisms is prevented.

### PK/PD relationship

For azithromycin the AUC/MIC is the major PK/PD parameter correlating best with the efficacy of azithromycin.

Following the assessment of studies conducted in children, the use of azithromycin is not recommended for the treatment of malaria, neither as monotherapy nor combined with chloroquine or artemisinin based drugs, as non-inferiority to anti-malarial drugs recommended in the treatment of uncomplicated malaria was not established.

### Mechanism of resistance:

Resistance to azithromycin may be inherent or acquired. There are three main mechanisms of resistance in bacteria: target site alteration, alteration in antibiotic transport and modification of the antibiotic.

Complete cross resistance exists among *Streptococcus pneumoniae*, beta-haemolytic streptococcus of group A, *Enterococcus faecalis* and *Staphylococcus aureus*, including methicillin resistant *S. aureus* (MRSA) to erythromycin, azithromycin, other macrolides and lincosamides.

### **Breakpoints**

EUCAST (European Committee on Antimicrobial Susceptibility Testing)

<b>Pathogens</b>	<b>susceptible (mg/l)</b>	<b>resistant (mg/l)</b>
<i>Staphylococcus</i> spp.	≤ 1	> 2
<i>Streptococcus</i> spp. (Group A, B, C, G)	≤ 0.25	> 0.5
<i>Streptococcus pneumoniae</i>	≤ 0.25	> 0.5
<i>Haemophilus influenzae</i>	≤ 0.12	> 4
<i>Moraxella catarrhalis</i>	≤ 0.5	> 0.5
<i>Neisseria gonorrhoeae</i>	≤ 0.25	> 0.5

### **Susceptibility:**

The prevalence of acquired resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert advice should be sought when the local prevalence of resistance is such that the utility of the agent in at least some types of infections is questionable.

Pathogens for which resistance may be a problem: prevalence of resistance is equal to or greater than 10% in at least one country in the European Union.

### **Table of susceptibility**

<b>Commonly susceptible species</b>
Aerobic Gram-negative microorganisms
<i>Haemophilus influenzae</i> *
<i>Moraxella catarrhalis</i> *
Other microorganisms
<i>Chlamydomphila pneumoniae</i>

<i>Chlamydia trachomatis</i>
<i>Legionella pneumophila</i>
<i>Mycobacterium avium</i>
<i>Mycoplasma pneumonia*</i>
<b>Species for which acquired resistance may be a problem</b>
Aerobic Gram-positive microorganisms
<i>Staphylococcus aureus*</i>
<i>Streptococcus agalactiae</i>
<i>Streptococcus pneumoniae*</i>
<i>Streptococcus pyogenes*</i>
Other microorganisms
<i>Ureaplasma urealyticum</i>
<b>Inherently resistant organisms</b>
Aerobic Gram-positive microorganisms
<i>Staphylococcus aureus</i> – methicillin resistant and erythromycin resistant strains
<i>Streptococcus pneumoniae</i> – penicillin resistant strains
Aerobic Gram-negative microorganisms
<i>Escherichia coli</i>
<i>Pseudomonas aeruginosa</i>
<i>Klebsiella</i> spp.
Anaerobic Gram-negative microorganisms
<i>Bacteroides fragilis</i> group

\* Clinical effectiveness is demonstrated by sensitive isolated organisms for approved clinical indications.

## 5.2 Pharmacokinetic properties

### *Absorption*

After oral administration the bioavailability of azithromycin is approximately 37%. Peak plasma levels are reached after 2-3 hours ( $C_{max}$  after a single dose of 500 mg orally was approximately 0.4 mg/l).

### *Distribution*

Kinetic studies have shown markedly higher azithromycin levels in tissue than in plasma (up to 50 times the maximum observed concentration in plasma) indicating that the active substance is heavily tissue bound (steady state distribution volume of approximately 31 l/kg). Concentrations in target tissues such as lung, tonsil, and prostate exceed the  $MIC_{90}$  for likely pathogens after a single dose of 500 mg.

In experimental *in vitro* and *in vivo* studies azithromycin accumulates in the phagocytes, freeing is stimulated by active phagocytosis. In animal studies this process appeared to contribute to the accumulation of azithromycin in the tissue.

In serum the protein binding of azithromycin is variable and depending on the serum concentration varies from 50% in 0.05 mg/l to 12% in 0.5 mg/l.



### *Excretion*

Plasma terminal elimination half-life closely reflects the tissue depletion half-life of 2 to 4 days. About 12% of an intravenously administered dose is excreted in the urine unchanged over a period of 3 days; the majority in the first 24 hours. Biliary excretion of azithromycin, predominantly in unchanged form, is a major route of elimination.

The identified metabolites (formed by N- and O- demethylising, by hydroxylising of the desosamine and aglycone rings, and by the splitting of the cladinose conjugate) are microbiologically inactive.

After a 5 day treatment slightly higher (29%) AUC values were seen in the elderly volunteers (>65 years of age) compared to the younger volunteers (< 45 years of age). However these differences are not regarded as clinically relevant; therefore a dose adjustment is not recommended.

### Pharmacokinetics in special populations

#### *Renal insufficiency*

Following a single oral dose of azithromycin 1 g, mean  $C_{max}$  and  $AUC_{0-120}$  increased by 5.1% and 4.2% respectively, in subjects with mild to moderate renal impairment (glomerular filtration rate of 10-80 ml/min) compared with normal renal function (GFR > 80 ml/min). In subjects with severe renal impairment, the mean  $C_{max}$  and  $AUC_{0-120}$  increased 61% and 33% respectively compared to normal.

#### *Hepatic insufficiency*

In patients with mild to moderate hepatic impairment, there is no evidence of a marked change in serum pharmacokinetics of azithromycin compared to normal hepatic function. In these patients, urinary recovery of azithromycin appears to increase perhaps to compensate for reduced hepatic clearance.

#### *Elderly*

The pharmacokinetics of azithromycin in elderly men was similar to that of young adults; however, in elderly women, although higher peak concentrations (increased by 30-50%) were observed, no significant accumulation occurred.

#### *Infants, toddlers, children and adolescents*

Pharmacokinetics have been studied in children aged 4 months – 15 years taking capsules, granules or suspension. At 10 mg/kg on day 1 followed by 5 mg/kg on days 2-5, the  $C_{max}$  achieved is slightly lower than adults with 224 ug/l in children aged 0.6-5 years and after 3 days dosing and 383 ug/l in those aged 6-15 years. The  $t_{1/2}$  of 36 h in the older children was within the expected range for adults.

### **5.3 Preclinical safety data**

In high-dose animal studies, giving active substance concentrations 40 fold higher than those expected in clinical practice, azithromycin has been noted to cause reversible phospholipidosis, generally without discernible toxicological consequences. There is no evidence that this is of relevance to the normal use of azithromycin in humans.

#### *Carcinogenic potential:*

Long-term studies in animals have not been performed to evaluate carcinogenic potential.

### *Mutagenic potential:*

Azithromycin has shown no mutagenic potential in standard laboratory tests: mouse lymphoma assay, human lymphocyte clastogenic assay, and mouse bone marrow clastogenic assay.

### *Reproductive toxicity:*

No teratogenic effects were observed in animal studies of embryotoxicity in mice and rats. In rats, azithromycin doses of 100 and 200 mg/kg bodyweight/day led to mild retardations in foetal ossification and in maternal weight gain. In peri-/postnatal studies in rats, mild retardations following treatment with 50 mg/kg/day azithromycin and above were observed.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Microcrystalline Cellulose  
Starch (Maize)  
Purified water  
Magnesium Stearate  
Purified Talc  
Sodium Starch glycolate  
Colloidal Silicon dioxide  
Croscarmellose Sodium  
Methylene chloride  
Isopropyl alcohol  
Titanium Dioxide as Wincoat Wt AQ-1001 White

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

36 Months

### **6.4 Special precautions for storage**

Store below 30° C.

### **6.5 Nature and contents of container**

Blister pack of 3 Tablets

Alu PVC blister of 3 Tablets. 1 such blisters are packed in one printed carton with pack insert.

## **6.6 Instructions for use and handling and disposal**

No special requirements.

## **7. MARKETING AUTHORISATION HOLDER UMEDICA LABORATORIES PVT. LTD.**

302, Dalamal House, 3rd Floor,  
J. Bajaj Road, Nariman Point,  
Mumbai 400 021, INDIA  
(Phone: +91 – 22 – 62455050, 40028503)  
Email : [exports@umedicalabs.com](mailto:exports@umedicalabs.com)

## **8. NUMBER(S) IN THE NATIONAL REGISTER OF FINISHED PHARMACEUTICAL PRODUCTS**

**Certificate No:** 07742/09555/NMR/2022

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

05/09/2022

## **10. DATE OF REVISION OF THE TEXT**

July 2023