

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE FINISHED PHARMACEUTICAL PRODUCT

Griseofulvin 500mg Tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION:

Label Claim: Each uncoated tablet contains
Griseofulvin BP...500 mg
Excipients q.s

3. PHARMACEUTICAL FORM: White to off white coloured round flat beveled uncoated tablet and break line on one side and plain on other side.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

The treatment of fungal infections of the skin, scalp, hair, or nails (Tinea barbae, Tinea capitis, Tinea corporis, Tinea cruris, Tinea pedis, Tinea unguium) where topical therapy is considered inappropriate, or the infection has proven refractory to topical therapy.

Oral administration of griseofulvin for systemic therapy of fungal infections enables newly formed keratin of the skin, hair, and nails to resist fungal attack. As the new keratin extends, the old infected keratin is shed.

Prior to therapy, the type of fungi responsible should be identified. The use of griseofulvin is not justified in the treatment of minor or trivial infections that will respond to topical therapy.

Before prescribing Griseofulvin Tablets, consideration should be given to national and/or local guidance on the appropriate use of antifungals.

4.2 Posology and method of administration

For oral administration.

Adults

The usual adult dose is 500mg to 1000mg daily. The dose should not be less than 10mg/kg bodyweight/ day. The dose may be administered as a single daily dose, or it may be administered twice daily. The twice daily dosing regimen may be more effective in those patients who respond poorly.

Elderly

No dosage adjustment is required in the elderly. Consideration should be given that such patients may also have a degree of hepatic impairment.

Children

The dose depends on the age and the weight of the child, and is usually 10mg/kg of the body weight per day. The dose usually given as two equal doses. Make sure you follow the instruction of Doctor.

4.3 Contraindications:

Griseofulvin is contraindicated in patients who have: -

- Hypersensitivity to griseofulvin or to any of the excipients, see section 6.1.
- Porphyria.
- Severe hepatic impairment.
- Systemic Lupus Erythematosus (SLE).
- Pregnancy, see section 4.6.
- Breastfeeding, see section 4.6.

4.4 Special warnings and precautions for use:

Griseofulvin is recommended after a high fat meal for increased absorption and minimising GI distress.

Griseofulvin is contraindicated in patients with severe hepatic impairment, see section 4.3. In patients with minor to moderate hepatic impairment, griseofulvin may cause further deterioration of hepatic function. Therefore care should be exercised with such patients, and it is recommended to perform regular periodic liver function tests, see section 4.8.

Griseofulvin is contraindicated in patients with Systemic Lupus Erythematosus (SLE), see section 4.3; griseofulvin has been reported to exacerbate the conditions, and care should be taken to exclude patients with pre-existing SLE from therapy.

Animal data, see section 5.3, indicates long term administration of high dose griseofulvin induces tumours in some species, but not others. The clinical relevance of this to man is unknown, but griseofulvin should not be used prophylactically.

Griseofulvin is a liver microsomal enzyme inducer and thus may impair the effectiveness of oral contraceptives. Therefore in women of child bearing age using oral contraception, additional barrier methods of contraception must be used during therapy and for 4 weeks following therapy cessation, see sections 4.5 and 4.6.

Griseofulvin causes chromosomal abnormalities in animals, see section 5.3. Therefore sexually active males should be cautioned to use an effective barrier method of contraception throughout therapy and for 6 months after therapy termination, see section 4.6.

A theoretical possibility of cross sensitivity in patients known to be allergic to penicillins exists, therefore caution should be exercised in administration of griseofulvin to such patients. It should be noted that such patients have been satisfactorily treated with griseofulvin without sequelae.

Patients should be cautioned to avoid excessive and unnecessary exposure to sunlight or U.V sources, including sunbeds, during griseofulvin therapy as photosensitivity reactions can occur, see section 4.8.

Consumption of alcohol in association with griseofulvin can result in an “Antabuse” type reaction, see section 4.5. Patients should be cautioned to avoid consumption of alcoholic beverages, and medicines containing alcohol, while undergoing griseofulvin therapy.

In patients undergoing long term griseofulvin therapy, i.e for tinea unguium, consideration should be given to periodic monitoring of blood chemistry, particularly for patients with pre-existing blood disorders, since griseofulvin may cause blood disorders, see section 4.8.

In common with any antibiotic, therapy with griseofulvin may result in the overgrowth of non-susceptible organisms, i.e bacteria or yeasts, or non-dermatophyte fungi, that are often cofactors in tinea infections, especially tinea pedis. Additional therapy is required to control or eradicate such organisms, as griseofulvin is ineffective.

Griseofulvin is not effective in infections due to *Candida albicans*, *Aspergillus* sp., *Malassezia furfur* (*Pityriasis versicolor*) and *Nocardia* sp. It has no antibacterial effects.

4.5 Interaction with other medicinal products and other forms of interaction:

Medicinal Products: Griseofulvin may depress plasma levels, and therefore the efficacy, of concomitantly administered medicinal products that are metabolised by cytochrome P450 3A4.

Interactions of Griseofulvin with other drugs:

Ciclosporin: concomitant administration may result in a reduction of ciclosporin plasma levels, necessitating a dosage adjustment. Plasma levels of ciclosporin should be monitored during griseofulvin therapy, and necessary dosage adjustments made.

Coumarin anticoagulants: the efficacy may be reduced, necessitating dosage adjustment. It is recommended that both prothrombin and INR are regularly monitored, for the duration of griseofulvin therapy, and for 8 days post therapy cessation.

Methadone: depression of methadone plasma levels may occur during griseofulvin therapy. Patients should be closely monitored for any loss of efficacy, or plasma levels of methadone be monitored, and corresponding dosage adjustments made.

Oral contraceptives: efficacy of oral contraception is reduced during griseofulvin therapy and for four weeks post therapy cessation. In view of the contraindication in pregnancy, see section 4.3, and of the possible sequelae of male patients fathering a child during therapy, all sexually active patients should use additional barrier contraception, such as condoms, throughout griseofulvin therapy, and for four weeks (female) and 6 months (male) post therapy cessation. See also sections 4.3, 4.4, 4.6, and 5.3 for additional information.

Interactions of other drugs with griseofulvin: Concurrent administration of other medicinal products that induce metabolising enzymes may result in a reduction of griseofulvin blood plasma levels and thus efficacy. The following drugs are known to have this effect:

Barbiturates, such as

- phenobarbitone
- Doxercalciferol
- Phenylbutazone
- Primidone

Other sedative and hypnotic drugs that induce metabolising enzymes.

Food: administration of griseofulvin after food, results in increased absorption, and thus higher plasma levels. This effect is enhanced if the meal contains high fat content. Administration after food is recommended, see section 4.2.

Alcohol: there are reports that griseofulvin enhances the central nervous system effects of alcohol. There are also reports that griseofulvin and alcohol use result in an “Antabuse” type reaction. Patients should be cautioned to avoid alcohol and all alcohol containing products while undergoing griseofulvin therapy, See also section 4.8.

4.6 Pregnancy and lactation

Pregnancy:

There are case reports of human foetal abnormalities associated with griseofulvin.

There are no adequate and well controlled studies in man, and inadequate epidemiological data. Griseofulvin has been shown to be teratogenic and embryotoxic in mice and rats. (see section 5.3).

Griseofulvin is suspected to cause serious birth defects when administered during pregnancy. Griseofulvin is contraindicated (see section 4.3) in pregnancy.

Women of childbearing potential have to use effective contraception during (and up to 4 weeks after) treatment (see section 4.5) in respect of effect on oral contraceptives, and contraceptive precautions.

Male-mediated effects on pregnancy Griseofulvin has been shown to induce chromosomal aberrations in animal spermatocytes (see section 5.3). Therefore men should take effective contraceptive precautions, i.e barrier contraception, to avoid fathering children for the duration of griseofulvin therapy, and for 6 months post therapy cessation.

Lactation:

It is unknown if griseofulvin is excreted in breast milk, but the possibility does exist. There is inadequate data on the safety of griseofulvin in breast feeding, and the potential risk to the infant cannot be assessed, therefore griseofulvin is contraindicated in breast feeding (see section 4.3).

4.7 Effects on ability to drive and use machines

Griseofulvin has no or negligible influence on the ability to drive and use machines. However, it may cause drowsiness, confusion dizziness, and impaired co-ordination, see section 4.8. Patients should therefore be cautioned not to drive or operate machines until they are sure they are not affected.

4.8 Undesirable effects

The following frequencies are used for the description of the occurrence of undesirable effects:

- Very common: $\geq 1 / 10$
- Common: $\geq 1 / 100, < 1 / 10$
- Uncommon: $\geq 1 / 1,000, < 1 / 100$
- Rare: $\geq 1 / 10,000, < 1 / 1,000$
- Very rare: $< 1 / 10,000$

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Headache and gastric discomfort are the most common effects on starting treatment, but usually disappear as treatment is continued.

Blood and lymphatic system disorder:

Rare: leucopenia, neutropenia, anaemia-these usually resolve on therapy cessation

Nervous system disorders:

Common: headache

Uncommon: impaired co-ordination, peripheral neuropathy, confusion, dizziness, drowsiness, insomnia, irritability.

Gastrointestinal disorders: Common: diarrhoea, vomiting, nausea, gastric discomfort

Uncommon: anorexia, taste sensation changes Skin and subcutaneous tissue disorders:

Uncommon: toxic epidermal necrolysis, erythema multiforme, photosensitivity on exposure to intense natural or artificial sunlight.

Rare: precipitation of Systemic Lupus Erythematosus, bullous reactions including Lyell's syndrome, urticarial reactions, skin rashes.

Hepatobiliary disorders:

Very rare: alteration in liver function tests, with elevation to more than three times upper normal limit, intrahepatic cholestasis, hepatitis.

4.8 Overdose

No case of overdose has been reported.

Symptoms

The likely symptoms of any overdose would be nausea, vomiting, headache, numbness and tingling, confusion, and vertigo. Urticaria or porphyria could occur.

Treatment:

There is no specific antidote to griseofulvin. Gastric lavage, or the induction of emesis may be of help, if ingestion is recent. Administration of activated charcoal may also be of use.

Treatment

There is no specific antidote to griseofulvin. Gastric lavage, or the induction of emesis may be of help, if ingestion is recent. Administration of activated charcoal may also be of use. Treatment should be symptomatic and supportive. Laboratory monitoring of haemopoetic, hepatic and nephritic parameters and electrolytes is recommended.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamics Properties

Pharmacotherapeutic group: Antifungals for systemic use

ATC code: D01BA01

Griseofulvin is an antifungal antibiotic that is active *in vivo* against common dermatophytes. The antifungal effect is manifested by binding to tubulin, at distinct binding sites, thus interfering with the microtubule function and causing inhibition of mitosis, and arresting cell division.

The inhibition of fungal mitosis leads to the production of multinucleate cells of characteristic morphology.

On entering the systemic circulation, griseofulvin binds to keratin in keratin precursor cells, thereby making them resistant to fungal infections. The drug only reaches the site of action when hair or skin is replaced by the keratingriseofulvin complex.

Griseofulvin then enters the dermatophyte through energy dependent transport processes and binds to the fungal microtubules, interfering with, and inhibiting mitosis, and the deposition of fungal cell walls.

Mycology:

Griseofulvin has antifungal activity against the following dermatophytes, although there is species and strain variability in susceptibility.

Trichophyton rubrum, *T. tonsurans*, *T. mentagrophytes*, *T. interdigitalis*, *T. verrucosum*, *T. megnini*, *T. gallinae*, *T. Crateriform*, *T. sulphureum* and *T. schoenleinii*.
Microsporum audouinii, *M. Canis*, *M. gypseum*.

Epidermophyton floccosum.

Griseofulvin has no activity against dermatophyte fungi of other genera, non-dermatophyte fungi, yeasts, gram positive bacteria, or gram negative bacteria. If any of these are cofactors in the pathology of infection, suitable additional therapy will be required for their eradication.

5.2 Pharmacokinetic properties

Absorption:

The absorption of griseofulvin from the gastrointestinal tract is variable and incomplete. On average, less than 50% of the oral dose is absorbed, but administration after a fatty meal, and a reduction in particle size will increase the rate and extent of the absorption.

Following oral administration there is a phase of rapid absorption, and thereafter a phase of slower prolonged absorption.

Peak plasma levels, 0.5 µg / ml-1.5 µg / ml after a 500 mg dose, and 1.5 µg / ml-3.0 µg / ml after a 1000 mg dose, are reached in 2-4 hours, and are maintained for some 10-20 hours. Griseofulvin exhibits linear pharmacokinetics.

Distribution:

The volume of distribution is about 0.7 L / Kg, and griseofulvin is ca 80 % bound to plasma proteins, predominantly serum albumin.

Griseofulvin crosses the placenta, and may be excreted in breast milk. There is selective deposition of griseofulvin in newly formed keratin of hair, skin, and nails, which gradually moves to the surface of these appendages.

Metabolism:

Griseofulvin undergoes metabolism to inactive metabolites, principally 6-desmethylgriseofulvin, or its glucuronide conjugate.

Excretion:

The terminal plasma half life ranges from 9.5-21 hours, with considerable intersubject variability. The majority of the dose, as 6-desmethylgriseofulvin or the glucuronide conjugate, and other metabolites is excreted in the urine, with less than 1% administered dose being excreted as unchanged griseofulvin. The remainder of the dose, principally as metabolites, is excreted in bile and faeces.

Renal insufficiency does not lead to accumulation.

5.3 PRECLINICAL SAFETY DATA

Griseofulvin can induce aneuploidy and meiotic delay in mouse oocytes following oral administration of high doses, i.e. 250mg/kg or greater. In addition, griseofulvin caused increases in numerical and structural chromosome aberrations in mouse spermatocytes at doses of 500mg/kg and above. Aneuploidy was observed at doses of 1500mg/kg.

Griseofulvin administered to rats and mice during pregnancy has been associated with foetotoxicity and foetal malformations. Long-term administration of high doses of

griseofulvin with food has been reported to induce hepatomas in mice and thyroid tumours in rats but not hamsters (see contraindications). The effects in mice may be due to a species specific effect on porphyrin metabolism.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Microcrystalline Cellulose, Maize Starch, Sodium Lauryl Sulfate, Povidone, Magnesium Stearate, Purified Talc.

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

36 Months

6.4 Special precautions for storage

Do not store above 30°C.

6.5 Nature and contents of container

Alu PVC Blister pack of 10 tablets and such 10 blisters are packed in a carton along with the pack insert.

6.6 Instructions for use and handling

Any unused product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Milan Laboratories (India) Pvt. Ltd.

303 & 304, Odyssey IT park,

Road No. 9, Opposite MIDC Office,

Wagle Estate, Thane -400604

India

E-mail: info@milanlabs.com

8. NUMBER(S) IN THE NATIONAL REGISTER OF FINISHED PHARMACEUTICAL PRODUCTS

06506/09012/NMR/2021

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

26.08.2021

10. DATE OF REVISION OF THE TEXT

Jul 2023