

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE FINISHED PHARMACEUTICAL PRODUCT

Cotrimoxazole Tablets BP 480 mg (MEDOPRIM 480)

2. QUALITATIVE AND QUANTITATIVE COMPOSITION:

Each uncoated tablet contains:

Sulfamethoxazole BP 400 mg and Trimethoprim BP 80 mg

For full list of excipients, see Section 6.1

3. PHARMACEUTICAL FORM

White circular flat, bevelled edge uncoated tablet with engraved as "COTRI" & "480" on one side and other side plain.

4. Clinical particulars

4.1 Therapeutic indications

Co-trimoxazole tablets are indicated in adults and children aged 6 years and over for the treatment of the following infections when owing to sensitive organisms:

- Treatment and prophylaxis of *Pneumocystis jirovecii* pneumonitis or "PJP"
- Treatment and prophylaxis of toxoplasmosis.
- Treatment of nocardiosis.

The following infections may be treated with co-trimoxazole where there is bacterial evidence of sensitivity to co-trimoxazole and good reason to prefer the combination of antibiotics in co-trimoxazole to a single antibiotic:

- Acute uncomplicated urinary tract infections
- Acute exacerbations of chronic bronchitis
- Acute otitis media

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

Posology

General Dosage Recommendations

Special Dosage Recommendations

(Standard dosage applies unless otherwise specified)

Where dosage is expressed as "tablets" this refers to the adult tablet, i.e. 80 mg Trimethoprim BP and 400 mg Sulfamethoxazole BP. If other formulations are to be used appropriate adjustment should be made.

Standard dosage recommendations for acute infections

Adults (>18 years): 2 tablets every 12 hours.

Paediatric population, children over 6 years old.

The standard dosage for children is equivalent to approximately 6 mg trimethoprim and 30 mg Sulfamethoxazole per kilogram body weight per day, given in two equally divided doses.

The schedules for children are according to the child's age and provided in the table below.

>12 to <18 years old	2 tablets every 12 hours
Children 6 to 12 years	1 tablet twice a day
Children under 6 years	Not recommended

Treatment should be continued until the patient has been symptom-free for two days; the majority will require treatment for at least 5 days. If clinical improvement is not evident after 7 days therapy, the patient should be reassessed.

As an alternative to standard dosage for acute uncomplicated lower urinary tract infections, short term therapy of 1 to 3 days duration has been shown to be effective.

The elderly:

See Special Warnings and Precautions for Use (section 4.4). Unless otherwise specified standard dosage applies.

Impaired hepatic function:

No data are available relating to dosage in patients with impaired hepatic function.

Impaired renal function:

Adults and children over 12 years:

Creatinine Clearance	Recommended Dosage
>30	2 tablets every 12 hours
15 to 30	1 tablet every 12 hours
<15	Not recommended

No information available for children aged 12 years and under with renal failure. See section 5.2 for the pharmacokinetics in the paediatric population with normal renal function of both components of Co-trimoxazole, TMP and SMZ.

Measurements of plasma concentration of sulfamethoxazole at intervals of 2–3 days are recommended in samples obtained 12 hours after administration of Co-trimoxazole. If the concentration of total sulfamethoxazole exceeds 150 mcg/ml then treatment should be interrupted until the value falls below 120 mcg/ml.

Pneumocystis jirovecii pneumonitis:

Treatment- Children (>12 to <18 years old) and adults (>18 years old):

A higher dosage is recommended, using 20 mg trimethoprim and 100 mg sulfamethoxazole per kg of body weight per day in two or more divided doses for two weeks. The aim is to obtain peak plasma or serum levels of trimethoprim of greater than or equal to 5 microgram/ml (verified in patients receiving 1-hour infusions of intravenous co-trimoxazole). (See 4.8 Undesirable Effects).

Prevention:

Adults (>18 years old): The following dose schedules may be used:

160 mg trimethoprim/800 mg sulfamethoxazole daily 7 days per week. 160 mg trimethoprim/800 mg sulfamethoxazole three times per week on alternate days.

320 mg trimethoprim/1600 mg sulfamethoxazole per day in two divided doses three times per week on alternate days.

The daily dose given on a treatment day approximates to 150 mg trimethoprim/m²/day and 750 mg sulfamethoxazole/m²/day. The total daily dose should not exceed 320 mg trimethoprim and 1600 mg sulfamethoxazole.

Children (>12 to <18 years old):

The standard dosage for children is equivalent to approximately 6 mg trimethoprim and 30 mg sulfamethoxazole per kg body weight per day, given in two equally divided doses. The following dose schedules may be used for the duration of the period at risk:

Age	Tablets
>12 to <18 years old	1 tablet every 12 hours, seven days per week
>12 to <18 years old	1 tablet every 12 hours, three times per week on
>12 to <18 years old	1 tablet every 12 hours, three times per week on
>12 to <18 years old	2 tablets once a day, three times per week on consecutive days

Children 6 – 12 years:

One tablet twice daily on 3 consecutive or alternate days per week or on 7 days per week.

Nocardiosis – Adults (>18 years old): There is no consensus on the most appropriate dosage. Adult doses of 6 to 8 tablets daily for up to 3 months have been used.

Toxoplasmosis: There is no consensus on the most appropriate dosage for the treatment or prophylaxis of this condition. The decision should be based on clinical experience. For

prophylaxis, however, the dosages suggested for prevention of *Pneumocystis jirovecii* pneumonitis may be appropriate.

Route of administration

Oral

It may be preferable to take co-trimoxazole with some food or drink to minimise the possibility of gastrointestinal disturbances.

4.3 Contraindications

- Hypersensitivity to the active substance(s), sulfonamides, trimethoprim, or to any of the excipients listed in section 6.1.
- Contra-indicated in patients showing marked liver parenchymal damage.
- Contra-indicated in severe renal insufficiency where repeated measurements of the plasma concentration cannot be performed.
- Co-trimoxazole should not be given to patients with a history of drug-induced immune thrombocytopenia with use of trimethoprim and/or sulfonamides.
- Co-trimoxazole should not be given to patients with acute porphyria.
- Co-trimoxazole should not be given to infants during the first 6 weeks of life.

4.4 Special warnings and precautions for use

Fatalities, although very rare, have occurred due to severe reactions including Stevens-Johnson syndrome, toxic epidermal necrolysis, fulminant hepatic necrosis, agranulocytosis, aplastic anaemia, other blood dyscrasias and hypersensitivity of the respiratory tract.

Life-threatening cutaneous reactions Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported with the use of Cotrimoxazole.

Patients should be advised of the signs and symptoms and monitored closely for skin reactions.

The highest risk for occurrence of SJS, TEN or DRESS is within the first weeks of treatment.

If symptoms or signs of SJS, TEN or DRESS (e.g. progressive skin rash often with blisters or mucosal lesions) are present, co-trimoxazole treatment should be discontinued (see 4.8 Undesirable Effects).

The best results in managing SJS, TEN and DRESS come from early diagnosis and immediate discontinuation of any suspect drug. Early withdrawal is associated with a better prognosis.

If the patient has developed SJS, TEN or DRESS with the use of Cotrimoxazole, Cotrimoxazole must not be re-started in this patient at any time.

The occurrence of a generalized febrile erythema associated with pustules, should raise the suspicion of acute generalised exanthematous pustulosis (AGEP) (see section 4.8); it requires cessation of treatment and contraindicates any new administration of Co-Trimoxazole alone or in combination with other drugs.

Respiratory toxicity

Very rare, severe cases of respiratory toxicity, sometimes progressing to Acute Respiratory Distress Syndrome (ARDS), have been reported during Cotrimoxazole treatment. The onset of pulmonary signs such as cough, fever, and dyspnoea in association with radiological signs of pulmonary infiltrates, and deterioration in pulmonary function may be preliminary signs of ARDS. In such circumstances, co-trimoxazole should be discontinued and appropriate treatment given.

Haemophagocytic lymphohistiocytosis (HLH)

Cases of HLH have been reported very rarely in patients treated with cotrimoxazole. HLH is a life-threatening syndrome of pathologic immune activation characterised by clinical signs and symptoms of an excessive systemic inflammation (e.g., fever, hepatosplenomegaly, hypertriglyceridaemia, hypofibrinogenaemia, high serum ferritin, cytopenias and haemophagocytosis). Patients who develop early manifestations of pathologic immune activation should be evaluated immediately. If diagnosis

of HLH is established, co-trimoxazole treatment should be discontinued.

Particular care is always advised when treating elderly patients because, as a group, they are more susceptible to adverse reactions and more likely to suffer serious effects as a result particularly when complicating conditions exist, e.g. impaired kidney and/or liver function and/or concomitant use of other drugs.

For patients with known renal impairment special measures should be adopted (see section 4.2).

An adequate urinary output should be maintained at all times. Evidence of crystalluria in vivo is rare, although sulfonamide crystals have been noted in cooled urine from treated patients. In patients suffering from malnutrition the risk may be increased.

Regular monthly blood counts are advisable when co-trimoxazole is given for long periods, or to folate deficient patients or to the elderly; since there exists a possibility of asymptomatic

changes in haematological laboratory indices due to lack of available folate. Supplementation with folinic acid may be considered during treatment but this should be initiated with caution due to possible interference with antimicrobial efficacy.

In glucose-6-phosphate-dehydrogenase (G-6-PD) deficient patients haemolysis may occur.

Co-trimoxazole should be given with caution to patients with severe allergy or bronchial asthma.

The administration of Co-trimoxazole to patients known or suspected to be at risk of acute porphyria should be avoided. Both trimethoprim and sulfonamides (although not specifically sulfamethoxazole) have been associated with clinical exacerbation of porphyria.

Trimethoprim has been noted to impair phenylalanine metabolism but this is of no significance in phenylketonuria patients on appropriate dietary restrictions.

Co-trimoxazole should not be used in the treatment of streptococcal pharyngitis due to group A beta-haemolytic streptococci; eradication of these organisms from the oropharynx is less effective than with penicillin.

Concomitant use of medicinal products known to cause hyperkalaemia with Co-trimoxazole may result in severe hyperkalaemia.

Close monitoring of serum potassium is warranted in patients at risk of hyperkalaemia.

Co-Trimoxazole has been associated with metabolic acidosis when other possible underlying causes have been excluded. Close monitoring is always advisable when metabolic acidosis is suspected.

Except under careful supervision Co-trimoxazole should not be given to patients with serious haematological disorders (see 4.8 Undesirable effects).

Co-trimoxazole has been given to patients receiving cytotoxic therapy with little or no additional effect on the bone marrow or peripheral blood.

The combination of antibiotics in Co-trimoxazole should only be used where, in the judgement of the physician, the benefits of treatment outweigh any possible risks; consideration should be

given to the use of a single effective antibacterial agent.

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'

4.5 Interaction with other medicinal products and other forms of interaction

Interaction with laboratory tests: trimethoprim may interfere with the estimation of serum/plasma creatinine when the alkaline picrate reaction is used. This may result in overestimation of serum/plasma creatinine of the order of 10%. The creatinine clearance is reduced: the renal tubular secretion of creatinine is decreased from 23% to 9% whilst the glomerular filtration remains unchanged.

Zidovudine: in some situations, concomitant treatment with zidovudine may increase the risk of haematological adverse reactions to co-trimoxazole. If concomitant treatment is necessary, consideration should be given to monitoring of haematological parameters.

Cyclosporin: reversible deterioration in renal function has been observed in patients treated with co-trimoxazole and cyclosporin following renal transplantation.

Rifampicin: concurrent use of rifampicin and Co-Trimoxazole results in a shortening of the plasma half-life of trimethoprim after a period of about one week. This is not thought to be of clinical significance.

When trimethoprim is administered simultaneously with drugs that form cations at physiological pH, and are also partly excreted by active renal secretion (e.g. procainamide, amantadine), there is the possibility of competitive inhibition of this process which may lead to an increase in plasma concentration of one or both of the drugs.

Diuretics (thiazides): in elderly patients concurrently receiving diuretics, mainly thiazides, there appears to be an increased risk of thrombocytopenia with or without purpura.

Pyrimethamine: occasional reports suggest that patients receiving pyrimethamine at doses in excess of 25 mg weekly may develop megaloblastic anaemia should co-trimoxazole be prescribed concurrently.

Warfarin: co-trimoxazole has been shown to potentiate the anticoagulant activity of warfarin via stereo-selective inhibition of its metabolism. Sulfamethoxazole may displace warfarin from plasma-albumin protein-binding sites in vitro. Careful control of the anticoagulant therapy during treatment with Co-Trimoxazole is advisable.

Phenytoin: co-trimoxazole prolongs the half-life of phenytoin and if co-administered could result in excessive phenytoin effect. Close monitoring of the patient's condition and serum phenytoin levels are advisable.

Digoxin: concomitant use of trimethoprim with digoxin has been shown to increase plasma digoxin levels in a proportion of elderly patients.

Methotrexate: co-trimoxazole may increase the free plasma levels of methotrexate. If Co-Trimoxazole is considered appropriate therapy in patients receiving other anti-folate drugs such as methotrexate, a folate supplement should be considered (see section 4.4).

Trimethoprim interferes with assays for serum methotrexate when dihydrofolate reductase from *Lactobacillus casei* is used in the assay. No interference occurs if methotrexate is measured by radioimmuno assay.

Lamivudine: administration of trimethoprim/sulfamethoxazole 80 mg/400 mg (co-trimoxazole) causes a 40% increase in lamivudine exposure because of the trimethoprim component. Lamivudine has no effect on the pharmacokinetics of trimethoprim or sulfamethoxazole.

Interaction with sulphonylurea hypoglycaemic agents is uncommon but potentiation has been reported.

Hyperkalaemia: caution should be exercised in patients taking any other drugs that can cause hyperkalaemia, for example ACE inhibitors, angiotensin receptor blockers and potassium-sparing diuretics such as spironolactone. Concomitant use of trimethoprim-sulfamethoxazole (co-trimoxazole) may result in clinically relevant hyperkalaemia.

Repaglinide: trimethoprim may increase the exposure of repaglinide which may result in hypoglycaemia.

Folic acid: folic acid supplementation has been shown to interfere with the antimicrobial efficacy of trimethoprim-sulfamethoxazole. This has been observed in *Pneumocystis jirovecii* pneumonia prophylaxis and treatment.

Contraceptives: oral contraceptive failures have been reported with antibiotics. The mechanism of this effect has not been elucidated. Women on treatment with antibiotics should temporarily use a barrier method in addition to the oral contraceptive, or choose another method of contraception.

Azathioprine: There are conflicting clinical reports of interactions between azathioprine and trimethoprim-sulfamethoxazole, resulting in serious haematological abnormalities.

4.6 Fertility, pregnancy and lactation

Pregnancy

Trimethoprim and sulfamethoxazole cross the placenta and their safety in pregnant women has not been established. Case-control studies have shown that there may be an association between exposure to folate antagonists and birth defects in humans.

Trimethoprim is a folate antagonist and, in animal studies, both agents have been shown to cause foetal abnormalities (see section 5.3). Co-Trimoxazole should not be used in pregnancy, particularly in the first trimester, unless clearly necessary. Folate supplementation should be considered if Co-Trimoxazole is used in pregnancy.

Sulfamethoxazole competes with bilirubin for binding to plasma albumin. As significantly maternally derived drug levels persist for several days in the newborn, there may be a risk of precipitating or exacerbating neonatal hyperbilirubinaemia, with an associated

theoretical risk of kernicterus, when Co-Trimoxazole is administered to the mother near the time of delivery. This theoretical risk is particularly relevant in infants at increased risk of hyperbilirubinaemia, such as those who are preterm and those with glucose-6-phosphate dehydrogenase deficiency.

Breast-feeding

The components of Co-Trimoxazole (trimethoprim and sulfamethoxazole) are excreted in breast milk. Administration of Co-Trimoxazole should be avoided in late pregnancy and in lactating mothers where the mother or infant has, or is at particular risk of developing, hyperbilirubinaemia. Additionally, administration of Co-Trimoxazole should be avoided in infants younger than eight weeks in view of the predisposition of young infants to hyperbilirubinaemia..

4.7 Effects on ability to drive and use machines

There have been no studies to investigate the effect of Co-Trimoxazole on driving performance or the ability to operate machinery. Further a detrimental effect on such activities cannot be predicted from the pharmacology of the drug. Nevertheless the clinical status of the patient and the adverse events profile of Co-Trimoxazole should be borne in mind when considering the patient's ability to operate machinery.

4.8 Undesirable effects

As co-trimoxazole contains trimethoprim and a sulphonamide the type and frequency of adverse reactions associated with such compounds are expected to be consistent with extensive historical experience.

Data from large published clinical trials were used to determine the frequency of very common to rare adverse events. Very rare adverse events were primarily determined from post-marketing experience data and therefore refer to reporting rate rather than a "true" frequency. In addition, adverse events may vary in their incidence depending on the indication.

The following convention has been used for the classification of adverse events in terms of frequency: Very common $\geq 1/10$, common $\geq 1/100$ and $< 1/10$, uncommon $\geq 1/1000$ and $< 1/100$, rare $\geq 1/10,000$ and $< 1/1000$, very rare $< 1/10,000$, not known - cannot be estimated from the available data.

System Organ Class	Frequency	Side effects
Infections and infestations	Common	Overgrowth fungal.
	Very rare	Pseudomembranous colitis

Blood and lymphatic system disorders	Veryrare	Leukopenia, neutropenia, thrombocytopenia, agranulocytosis, anaemia megaloblastic, aplastic anaemia, haemolytic anaemia, methaemoglobinemia, eosinophilia, purpura, haemolysis in certain susceptible G-6-PD
Immune system disorders	Veryrare	Serum sickness, anaphylactic reactions, allergic myocarditis, hypersensitivity vasculitis resembling Hensch-Schoenlein purpura, periarteritis nodosa, systemic lupus erythematosus. Severe hypersensitivity reactions associated with PJP*, rash, pyrexia, neutropenia, thrombocytopenia, hepatic enzyme increased, hyperkalaemia,
Metabolism and nutrition disorders	Verycommon	Hyperkalaemia.
	Veryrare	Hypoglycaemia, hyponatraemia, decreased appetite, metabolic acidosis
Psychiatric disorders	Veryrare	Depression, hallucination.
	Not known	Psychotic disorder.
Nervous system disorders	Common	Headache.
	Veryrare	Meningitis aseptic *, seizure, neuropathy peripheral, ataxia, dizziness.
Ear and Labrynth disorders	Veryrare	Vertigo, tinnitus
Eyedisorders	Veryrare	Uveitis.
Respiratory, thoracic and mediastinal	Veryrare	Cough*, dyspnoea*, lung infiltration*.
Gastrointestinal disorders	Common	Nausea, diarrhoea.
	Uncommon	Vomiting.
	Veryrare	Glossitis, stomatitis, pancreatitis.
Hepatobiliary disorders	Veryrare	Jaundice cholestatic *, hepatic necrosis*. Transaminases increased, blood bilirubin increased.

Skin and subcutaneous tissue disorders*	Common	Rash.
	Very rare	Photosensitivity reaction, dermatitis exfoliative, angioedema, fixed drug eruption, erythema multiforme, Stevens-Johnson syndrome (SJS)*, toxic epidermal necrolysis (TEN)*.
Musculoskeletal and connective tissue disorders	Very rare	Arthralgia, myalgia.
Renal and urinary disorders	Very rare	Renal impairment (sometimes reported as renal failure), tubulointerstitial nephritis and uveitis syndrome, renal tubular acidosis.

*see description of selected adverse reactions

Description of selected

adverse reactions *Aseptic meningitis*

Aseptic meningitis was rapidly reversible on withdrawal of the drug, but recurred in a number of cases on re-exposure to either co-trimoxazole or to trimethoprim alone.

Pulmonary hypersensitivity reactions

Cough, dyspnoea and lung infiltration may be early indicators of respiratory hypersensitivity which, while very rare, has been fatal.

Hepatobiliary disorders

Jaundice cholestatic and hepatic necrosis may be fatal.

Skin and subcutaneous tissue disorders

Common: Skin rashes

Very rare: Photosensitivity, exfoliative dermatitis, fixed drug eruption, erythema multiforme.

Severe cutaneous adverse reactions (SCARs)

Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) have been reported

*Effects associated with *Pneumocystis jirovecii* Pneumonitis (PJP) management*

Very rare: Severe hypersensitivity reactions, rash, pyrexia, neutropenia, thrombocytopenia, hepatic enzyme increased, hyperkalaemia, hyponatraemia, rhabdomyolysis.

At the high dosages used for PJP management severe hypersensitivity reactions have been reported, necessitating cessation of therapy. Severe hypersensitivity reactions have been reported in PJP patients on re-exposure to co-trimoxazole, sometimes after a dosage interval of a few days. Rhabdomyolysis has been reported in HIV positive patients receiving co-trimoxazole for prophylaxis or treatment of PJP.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme (www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

4.9 Overdose

Symptoms:

Nausea, vomiting, dizziness and confusion are likely signs/symptoms of over dosage. Bone marrow depression has been reported in acute trimethoprim over dosage.

Treatment:

If vomiting has not occurred, induction of vomiting may be desirable. Gastric lavage may be useful, though absorption from the gastrointestinal tract is normally very rapid and complete within approximately two hours. This may not be the case in gross over dosage. Dependent on the status of renal function administration of fluids is recommended if urine output is low.

Both trimethoprim and active Sulfamethoxazole are moderately dialysable by haemodialysis. Peritoneal dialysis is not effective.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Combinations of sulphonamides and trimethoprim, incl. derivatives, **ATC code:** J01EE01.

Mechanism of Action

Sulfamethoxazole competitively inhibits the utilisation of para-aminobenzoic acid in the synthesis of dihydrofolate by the bacterial cell resulting in bacteriostasis. Trimethoprim reversibly inhibits bacterial dihydrofolate reductase (DHFR), an enzyme active in the folate metabolic pathway converting dihydrofolate to tetrahydrofolate. Depending on the condition the effect may be bactericidal. Thus trimethoprim and sulfamethoxazole block two consecutive steps in the biosynthesis of purines and therefore nucleic acids essential to many bacteria. This action produces marked potentiation of activity in vitro between the two agents.

Trimethoprim reversibly inhibits bacterial dihydrofolate reductase (DHFR), an enzyme active in the folate metabolic pathway converting dihydrofolate to tetrahydrofolate. Depending on the conditions the effect may be bactericidal. Thus trimethoprim and sulfamethoxazole block two consecutive steps in the biosynthesis of purines and therefore nucleic acids essential to many

bacteria. This action produces marked potentiation of activity in vitro between the two agents.

Trimethoprim binds to plasmodial DHFR but less tightly than to the bacterial enzyme. Its affinity for mammalian DHFR is some 50,000 times less than for the corresponding bacterial enzyme.

Mechanism of resistance

In vitro studies have shown that bacterial resistance can develop more slowly with both sulfamethoxazole and trimethoprim in combination than with either sulfamethoxazole or trimethoprim alone.

Resistance to sulfamethoxazole may occur by different mechanisms. Bacterial mutations cause an increase in the concentration of PABA and thereby out-compete with sulfamethoxazole resulting in a reduction of the inhibitory effect on dihydropteroate synthetase enzyme. Another resistance mechanism is plasmid-mediated and results from production of an altered dihydropteroate synthetase enzyme, with reduced affinity for sulfamethoxazole compared to the wild-type enzyme.

Resistance to trimethoprim occurs through a plasmid-mediated mutation which results in production of an altered dihydrofolate reductase enzyme having a reduced affinity for trimethoprim compared to the wild-type enzyme.

Trimethoprim binds to plasmodial DHFR but less tightly than to bacterial enzyme. Its affinity for mammalian DHFR is some 50,000 times less than for the corresponding bacterial enzyme.

Many common pathogenic bacteria are susceptible in vitro to trimethoprim and sulfamethoxazole at concentrations well below those reached in blood, tissue fluids and urine after administration of recommended doses. In common with other antibiotics, however, in vitro activity does not necessarily imply that clinical efficacy has been demonstrated and it must be noted that satisfactory susceptibility testing is achieved only with recommended media free from inhibitory substances, especially thymidine and thymine.

Susceptibility testing breakpoints

EUCAST

Enterobacteriaceae: $S \leq 2$ $R > 4$

S. maltophilia: $S \leq 4$ $R > 4$

Acinetobacter: $S \leq 2$ $R > 4$

Staphylococcus: $S \leq 2$ $R > 4$

Enterococcus: $S \leq 0.032$ $R > 1$

Streptococcus ABCG: $S \leq 1$ $R > 2$

Streptococcus pneumoniae: $S \leq 1$ $R > 2$

Haemophilus influenzae: $S \leq 0.5$ $R > 1$

Moraxella catarrhalis: $S \leq 0.5$ $R > 1$

Pseudomonas aeruginosa and other non-enterobacteriaceae: $S \leq 2^*$ $R > 4^*$

S=susceptible, R=resistant. *These are CLSI breakpoint values since no EUCAST breakpoints are currently available for these organisms.

MEDOPHARM PRIVATE LIMITED, INDIA

Trimethoprim: sulfamethoxazole in the ratio 1:19. Breakpoints are expressed as trimethoprim concentration.

Antibacterial Spectrum

The prevalence of resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert advice should be sought when the local prevalence of resistance is such that the utility of the agent in at least some types of infections is questionable. This information gives only an approximate guidance on probabilities whether microorganisms will be susceptible to trimethoprim/sulfamethoxazole or not.

Trimethoprim/sulfamethoxazole susceptibility against a number of bacteria are shown in the table below:

Commonly susceptible species:
Gram-positive aerobes: Staphylococcus aureus Staphylococcus saprophyticus Streptococcus pyogenes
Gram-negative aerobes: Enterobacter cloacae Haemophilus influenzae Klebsiella oxytoca Moraxella catarrhalis Salmonella spp. Stenotrophomonas maltophilia Yersinia spp.
Species for which acquired resistance may be a problem:
Gram-positive aerobes: Enterococcus faecalis Enterococcus faecium Nocardia spp. Staphylococcus epidermidis Streptococcus pneumoniae

<p>Gram-negative aerobes: Citrobacter spp. Enterobacter aerogenes Escherichia coli Klebsiella pneumoniae Klebsiella pneumonia Proteus mirabilis Proteus vulgaris Providencia spp. Serratia marcescens</p>
<p>Inherently resistant organisms:</p>
<p>Gram-negative aerobes: Pseudomonas aeruginosa Shigella spp. Vibrio cholera</p>

5.2 Pharmacokinetic properties

Absorption

After oral administration trimethoprim and sulfamethoxazole are rapidly and nearly completely absorbed. The presence of food does not appear to delay absorption. Peak levels in the blood occur between one and four hours after ingestion and the level attained is dose related. Effective levels persist in the blood for up to 24 hours after a therapeutic dose. Steady state levels in adults are reached after dosing for 2-3 days. Neither component has an appreciable effect on the concentrations achieved in the blood by the other.

Distribution

Approximately 50% of trimethoprim in the plasma is protein bound. Tissue levels of trimethoprim are generally higher than corresponding plasma levels, the lungs and kidneys showing especially high concentrations. Trimethoprim concentrations exceed those in plasma in the case of bile, prostatic fluid and tissue, saliva, sputum and vaginal secretions. Levels in the aqueous humor, breast milk, cerebrospinal fluid, middle ear fluid, synovial fluid and tissue (intestinal) fluid are adequate for antibacterial activity. Trimethoprim passes into amniotic fluid and foetal tissues reaching concentrations approximating those of maternal serum.

Approximately 66% of sulfamethoxazole in the plasma is protein bound. The concentration of active sulfamethoxazole in amniotic fluid, aqueous humour, bile, cerebrospinal fluid, middle ear fluid, sputum, synovial fluid and tissue (interstitial) fluids is of the order of 20 to 50% of the plasma concentration

Biotransformation

Renal excretion of intact sulfamethoxazole accounts for 15-30% of the dose. This drug is more extensively metabolised than trimethoprim, via acetylation, oxidation or glucuronidation. Over a 72 hour period, approximately 85% of the dose can be accounted for in the urine as unchanged drug plus the major (N4-acetylated) metabolite.

Elimination

The half-life of trimethoprim in man is in the range 8.6 to 17 hours in the presence of normal renal function. It is increased by a factor of 1.5 to 3.0 when the creatinine clearance is less than 10 ml/minute. There appears to be no significant difference in elderly patients compared with young patients.

The principal route of excretion of trimethoprim is renal and approximately 50% of the dose is excreted in the urine within 24 hours as unchanged drug. Several metabolites have been identified in the urine. Urinary concentrations of trimethoprim vary widely.

The half-life of sulfamethoxazole in man is approximately 9 to 11 hours in the presence of normal renal function.

There is no change in the half-life of active sulfamethoxazole with a reduction in renal function but there is prolongation of the half-life of the major, acetylated metabolite when the creatinine clearance is below 25 ml/minute.

The principal route of excretion of sulfamethoxazole is renal; between 15% and 30% of the dose recovered in the urine is in the active form.

The pharmacokinetics in the paediatric population with normal renal function of both components of Cotrimoxazole, TMP and SMZ are age dependent. Elimination of TMP-SMZ is reduced in neonates, during the first two months of life, thereafter both TMP and SMZ show a higher elimination with a higher body clearance and a shorter elimination half-life. The differences are most prominent in young infants (> 1.7 months up to 24 months) and decrease with increasing age, as compared to young children (1 year up to 3.6 years), children (7.5 years and < 10 years) and adults.

Special patient population

Renal impairment

The elimination half-life of trimethoprim is increased by a factor of 1.5-3.0 when the creatinine clearance

is less than 10mL/minute. When the creatinine clearance falls below 30 mL/min the dosage of Co-Trimoxazole should be reduced.

Hepatic impairment

Caution should be exercised when treating patients with severe hepatic parenchymal damage as there may be changes in the absorption and biotransformation of trimethoprim and sulfamethoxazole.

Elderly patients

In elderly patients, a slight reduction in renal clearance of Sulfamethoxazole but not trimethoprim has been observed.

Paediatric population

See special dosage regimen.

5.3 Preclinical safety data

Reproductive toxicology: At doses in excess of recommended human therapeutic dose, trimethoprim and sulfamethoxazole have been reported to cause cleft palate and other foetal abnormalities in rats, findings typical of a folate antagonist. Effects with trimethoprim were preventable by administration of dietary folate. In rabbits, foetal loss was seen at doses of trimethoprim in excess of human therapeutic doses

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

- Maize starch BP
- Povidone K 30 BP
- Sodium lauryl sulphate BP
- Purified water BP
- Pre-gelatinized Starch BP
- Talc USP
- Sodium Starch Glycolate BP
- Colloidal Silicon Dioxide BP
- Magnesium Stearate BP

6.2 Incompatibilities

Not applicable

6.3 Shelf life

36 months

6.4 Special precautions for storage

Store in a dry place below 30°C. Protect from light.

6.5 Nature and contents of container

Presentation: Co – Trimoxazole tablets BP 480mg is available as 10 x 10's, 100 x 10's Blister pack and 1000's Bulk pack..

Primary Container (s):

Co – Trimoxazole tablets BP 480mg is available as Blister pack.

10x10's & 100x 10's - Each blister contains 10 tablets

Secondary packing:

Such blisters are packed in cartons of GSM 300, made of ITC cyber XL board with aqua varnish. Carton is printed in Multicolor.

Leaflet: leaflet made with 70 GSM Map Lithopaper.

Outer Container:

Such cartons are packed in Export Worthy 5/7 Ply Shippers. These shippers are labelled with product name and relevant batch details and sealed with BOPP tape. Shippers are then strapped with Polypropylene tapes.

Transportation: Should be transported with precautions.

The Cautions Like- This Side Up

- Not For Loose Handling
- Protect from Water
- Avoid Vigorous Transportation Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

None

7. MARKETING AUTHORIZATIONHOLDER

Name and Permanent address of the Marketing authorization holder:

Medopharm PrivateLimited

“MEDO HOUSE”

25, Puliur II Main road, Trustpuram, Chennai-600 024, Tamil Nadu, India.

PH: +91 44-30149992/30149955

Fax: 260211 286283

Manufacturing Site address:

Medopharm Private Limited (Unit II),

No. 50, Kayarambedu Village, Guduvanchery- 603 202, Tamilnadu, India.

8. NUMBER (S) IN THE NATIONAL REGISTER OF FINISHED PHARMACEUTICAL PRODUCTS

08114/09875/NMR/2022

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION:

13.11.2022

10. DATE OF REVISION OF THE TEXT

11.07.2023

11. REFERENCE

- <https://www.medicines.org.uk/emc/product/6999/smpc>