

## **SUMMARY OF PRODUCT CHARACTERISTICS**

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. Name of the medicinal product

Levo-Denk 500

### 2. Qualitative and quantitative composition

1 film-coated tablet contains 500 mg levofloxacin, equivalent to 512.46 mg levofloxacin hemihydrate.

For a full list of excipients, see section 6.1.

### 3. Pharmaceutical form

Film-coated tablet.

Pink, oblong, film-coated tablet with a breaking line on both sides.

The film-coated tablet can be divided into equal halves.

### 4. Clinical particulars

#### 4.1 Therapeutic indications

Levo-Denk 500 is indicated in adults for the treatment of the following infections (see sections 4.4 and 5.1):

- Acute bacterial sinusitis
- Acute exacerbations of chronic bronchitis
- Community-acquired pneumonia
- Complicated skin and soft tissue infections

For the above-mentioned infections Levo-Denk 500 should be used only when it is considered inappropriate to use antibacterial agents that are commonly recommended for the initial treatment of these infections.

- Pyelonephritis and complicated urinary tract infections (see section 4.4)
- Chronic bacterial prostatitis
- Uncomplicated cystitis (see section 4.4)
- Inhalation Anthrax: postexposure prophylaxis and curative treatment (see section 4.4)

Levo-Denk 500 may also be used to complete a course of therapy in patients who have shown improvement during initial treatment with intravenous levofloxacin.

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

#### 4.2 Posology and method of administration

Levo-Denk 500 tablets are administered once or twice daily. The dosage depends on the type and severity of the infection and the susceptibility of the presumed causative pathogen.

Levo-Denk 500 tablets may also be used to complete a course of therapy in patients who have shown improvement during initial treatment with intravenous levofloxacin; given the bioequivalence of the parenteral and oral forms, the same dosage can be used.

#### Posology

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

The following dose recommendations can be given for Levo-Denk 500:

*Dosage in patients with normal renal function (creatinine clearance >50 ml/min)*

<b>Indication</b>	<b>Daily dose regimen (according to severity)</b>	<b>Duration of treatment (according to severity)</b>
Acute bacterial sinusitis	500 mg once daily	10 - 14 days
Acute bacterial exacerbations of chronic bronchitis	500 mg once daily	7 - 10 days
Community-acquired pneumonia	500 mg once or twice daily	7 - 14 days
Pyelonephritis	500 mg once daily	7 - 10 days
Complicated urinary tract infections	500 mg once daily	7 - 14 days
Uncomplicated cystitis	250 mg once daily	3 days
Chronic bacterial prostatitis	500 mg once daily	28 days
Complicated skin and soft tissue infections	500 mg once or twice daily	7 - 14 days
Inhalation Anthrax	500 mg once daily	8 weeks

### Special populations

*Impaired renal function (creatinine clearance ≤50 ml/min)*

	<b>Dose regimen</b>		
	<b>250 mg/24 h</b>	<b>500 mg/24 h</b>	<b>500 mg/12 h</b>
<b>Creatinine clearance</b>	<i>first dose: 250 mg</i>	<i>first dose: 500 mg</i>	<i>first dose: 500 mg</i>
50-20 ml/min	<i>then: 125 mg/24 h</i>	<i>then: 250 mg/24 h</i>	<i>then: 250 mg/12 h</i>
19-10 ml/min	<i>then: 125 mg/48 h</i>	<i>then: 125 mg/24 h</i>	<i>then: 125 mg/12 h</i>
<10 ml/min (including haemodialysis and CAPD) <sup>1</sup>	<i>then: 125 mg/48 h</i>	<i>then: 125 mg/24 h</i>	<i>then: 125 mg/24 h</i>

<sup>1</sup>No additional doses are required after haemodialysis or continuous ambulatory peritoneal dialysis (CAPD).

### *Impaired liver function*

No adjustment of dose is required since levofloxacin is not metabolised to any relevant extent by the liver and is mainly excreted by the kidneys.

### *Elderly population*

No adjustment of dose is required in the elderly, other than that imposed by consideration of renal function (see section 4.4 “Tendinitis and tendon rupture” and “QT interval prolongation”).

### *Paediatric population*

Levo-Denk 500 is contraindicated in children and growing adolescents (see section 4.3).

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

#### Method of administration

Levo-Denk 500 tablets should be swallowed without crushing and with sufficient amount of liquid. They may be divided at the score line to adapt the dose. The tablets may be taken during meals or between meals. Levo-Denk 500 tablets should be taken at least two hours before or after iron salts, zinc salts, magnesium- or aluminium-containing antacids, or didanosine (*only didanosine formulations with aluminium or magnesium containing buffering agents*), and sucralfate administration, since reduction of absorption can occur (see section 4.5).

### **4.3 Contraindications**

Levofloxacin tablets must not be used:

- in patients hypersensitive to levofloxacin or other quinolones or any of the excipients listed in section 6.1,
- in patients with epilepsy,
- in patients with history of tendon disorders related to fluoroquinolone administration,
- in children or growing adolescents,
- during pregnancy,
- in breast-feeding women.

### **4.4 Special warnings and precautions for use**

Methicillin-resistant *S. aureus* are very likely to possess co-resistance to fluoroquinolones, including levofloxacin. Therefore levofloxacin is not recommended for the treatment of known or suspected MRSA infections unless laboratory results have confirmed susceptibility of the organism to levofloxacin (and commonly recommended antibacterial agents for the treatment of MRSA-infections are considered inappropriate).

Levofloxacin may be used in the treatment of Acute Bacterial Sinusitis and Acute Exacerbation of Chronic Bronchitis when these infections have been adequately diagnosed.

Resistance to fluoroquinolones of *E. coli* – the most common pathogen involved in urinary tract infections – varies across the European Union. Prescribers are advised to take into account the local prevalence of resistance in *E. coli* to fluoroquinolones.

Inhalation Anthrax: Use in humans is based on *in vitro Bacillus anthracis* susceptibility data and on animal experimental data together with limited human data. Treating physicians should refer to national and/or international consensus documents regarding the treatment of anthrax.

#### *Tendinitis and tendon rupture*

Tendinitis may rarely occur. It most frequently involves the Achilles tendon and may lead to tendon rupture. Tendinitis and tendon rupture, sometimes bilateral, may occur within 48 hours of starting treatment with levofloxacin and have been reported up to several months after discontinuation of treatment. The risk of tendinitis and tendon rupture is increased in patients aged over 60 years, in patients receiving daily doses of 1000 mg and in patients using corticosteroids. The daily dose should be adjusted in elderly patients based on creatinine clearance (see section 4.2). Close monitoring of these patients is therefore necessary if they are prescribed levofloxacin. All patients should consult their physician if they experience symptoms of tendinitis. If tendinitis is suspected, treatment with levofloxacin must be halted immediately, and appropriate treatment (e.g. immobilisation) must be initiated for the affected tendon (see sections 4.3 and 4.8).

#### *Clostridium difficile-associated disease*

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

Diarrhoea, particularly if severe, persistent and/or bloody, during or after treatment with levofloxacin (including several weeks after treatment), may be symptomatic of *Clostridium difficile*-associated disease (CDAD). CDAD may range in severity from mild to life threatening, the most severe form of which is pseudomembranous colitis (see section 4.8). It is therefore important to consider this diagnosis in patients who develop serious diarrhoea during or after treatment with levofloxacin. If CDAD is suspected or confirmed, levofloxacin should be stopped immediately and appropriate treatment initiated without delay. Anti-peristaltic medicinal products are contraindicated in this clinical situation.

#### *Patients predisposed to seizures*

Quinolones may lower the seizure threshold and may trigger seizures. Levofloxacin is contraindicated in patients with a history of epilepsy (see section 4.3) and, as with other quinolones, should be used with extreme caution in patients predisposed to seizures or concomitant treatment with active substances that lower the cerebral seizure threshold, such as theophylline (see section 4.5). In case of convulsive seizures (see section 4.8), treatment with levofloxacin should be discontinued.

#### *Patients with G-6-phosphate dehydrogenase deficiency*

Patients with latent or actual defects in glucose-6-phosphate dehydrogenase activity may be prone to haemolytic reactions when treated with quinolone antibacterial agents. Therefore, if levofloxacin has to be used in these patients, potential occurrence of haemolysis should be monitored.

#### *Patients with renal impairment*

Since levofloxacin is excreted mainly by the kidneys, the dose of Levo-Denk 500 should be adjusted in patients with renal impairment (see section 4.2).

#### *Hypersensitivity reactions*

Levofloxacin can cause serious, potentially fatal hypersensitivity reactions (e.g. angioedema up to anaphylactic shock), occasionally following the initial dose (see section 4.8). Patients should discontinue treatment immediately and contact their physician or an emergency physician, who will initiate appropriate emergency measures.

#### *Severe bullous reactions*

Cases of severe bullous skin reactions such as Stevens-Johnson syndrome or toxic epidermal necrolysis have been reported with levofloxacin (see section 4.8). Patients should be advised to contact their doctor immediately prior to continuing treatment if skin and/or mucosal reactions occur.

#### *Dysglycaemia*

As with all quinolones, disturbances in blood glucose, including both hypoglycaemia and hyperglycaemia have been reported, usually in diabetic patients receiving concomitant treatment with an oral hypoglycaemic agent (e.g., glibenclamide) or with insulin. Cases of hypoglycaemic coma have been reported. In diabetic patients, careful monitoring of blood glucose is recommended (see section 4.8).

#### *Prevention of photosensitisation*

Photosensitisation has been reported with levofloxacin (see section 4.8). It is recommended that patients should not expose themselves unnecessarily to strong sunlight or to artificial UV rays (e.g. sunray lamp, solarium), during treatment and for 48 hours following treatment discontinuation in

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

order to prevent photosensitisation.

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

#### *Patients treated with Vitamin K antagonists*

Due to possible increase in coagulation tests (PT/INR) and/or bleeding in patients treated with levofloxacin in combination with a vitamin K antagonist (e.g. warfarin), coagulation tests should be monitored when these drugs are given concomitantly (see section 4.5).

#### *Psychotic reactions*

Psychotic reactions have been reported in patients receiving quinolones, including levofloxacin. In very rare cases these have progressed to suicidal thoughts and self-endangering behaviour-sometimes after only a single dose of levofloxacin (see section 4.8). In the event that the patient develops these reactions, levofloxacin should be discontinued and appropriate measures instituted. Caution is recommended if levofloxacin is to be used in psychotic patients or in patients with history of psychiatric disease.

#### *QT interval prolongation*

Caution should be taken when using fluoroquinolones, including levofloxacin, in patients with known risk factors for prolongation of the QT interval such as, for example:

- congenital long QT syndrome
- concomitant use of drugs that are known to prolong the QT interval (e.g. Class IA and III antiarrhythmics, tricyclic antidepressants, macrolides, antipsychotics)
- uncorrected electrolyte imbalance (e.g. hypokalemia, hypomagnesemia)
- cardiac disease (e.g. heart failure, myocardial infarction, bradycardia).

Elderly patients and women may be more sensitive to QTc-prolonging medications. Therefore, caution should be taken when using fluoroquinolones, including levofloxacin, in these populations. (See sections 4.2 *Elderly*, 4.5, 4.8, and 4.9).

#### *Peripheral neuropathy*

Peripheral sensory neuropathy and peripheral sensory motor neuropathy have been reported in patients receiving fluoroquinolones, including levofloxacin, which can be rapid in its onset (see section 4.8). Levofloxacin should be discontinued if the patient experiences symptoms of neuropathy in order to prevent the development of an irreversible condition.

#### *Hepatobiliary disorders*

Cases of hepatic necrosis up to fatal hepatic failure have been reported with levofloxacin, primarily in patients with severe underlying diseases, e.g. sepsis (see section 4.8). Patients should be advised to stop treatment and contact their doctor if signs and symptoms of hepatic disease develop such as anorexia, jaundice, dark urine, pruritus or tender abdomen.

#### *Exacerbation of myasthenia gravis*

Fluoroquinolones, including levofloxacin, have neuromuscular blocking activity and may exacerbate muscle weakness in patients with myasthenia gravis. Postmarketing serious adverse reactions, including deaths and the requirement for respiratory support, have been associated with fluoroquinolone use in patients with myasthenia gravis. Levofloxacin is not recommended in patients with a known history of myasthenia gravis.

#### *Vision disorders*

If vision becomes impaired or any effects on the eyes are experienced, an eye specialist should be consulted immediately (see sections 4.7 and 4.8).

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

### *Superinfection*

The use of levofloxacin, especially if prolonged, may result in overgrowth of non-susceptible organisms. If superinfection occurs during therapy, appropriate measures should be taken.

### *Interference with laboratory tests*

In patients treated with levofloxacin, determination of opiates in urine may give false-positive results. It may be necessary to confirm positive opiate screens by more specific method.

Levofloxacin may inhibit the growth of *Mycobacterium tuberculosis* and, therefore, may give false-negative results in the bacteriological diagnosis of tuberculosis.

### *Further precautionary measures*

Patients suffering from the rare hereditary galactose intolerance, lactase deficiency or glucose galactose malabsorption should not take Levo-Denk 500.

## **4.5 Interactions with other medicinal products and other forms of interaction**

### Effect of other medicinal products on Levo-Denk 500

#### *Iron salts, zinc salts, magnesium- or aluminium-containing antacids, didanosine*

Levofloxacin absorption is significantly reduced when iron salts, or magnesium- or aluminium-containing antacids, or didanosine (*only didanosine formulations with aluminium or magnesium containing buffering agents*) are administered concomitantly with Levo-Denk 500 tablets.

Concurrent administration of fluoroquinolones with multi-vitamins containing zinc appears to reduce their oral absorption. It is recommended that preparations containing divalent or trivalent cations such as iron salts, zinc salts or magnesium- or aluminium-containing antacids, or didanosine (*only didanosine formulations with aluminium or magnesium containing buffering agents*) should not be taken 2 hours before or after Levo-Denk 500 tablet administration (see section 4.2). Calcium salts have a minimal effect on the oral absorption of levofloxacin.

#### *Sucralfate*

The bioavailability of Levo-Denk 500 tablets is significantly reduced when administered together with sucralfate. If the patient is to receive both sucralfate and Levo-Denk 500, it is best to administer sucralfate 2 hours after the Levo-Denk 500 tablet administration (see section 4.2).

#### *Theophylline, fenbufen or similar non-steroidal anti-inflammatory drugs*

No pharmacokinetic interactions of levofloxacin were found with theophylline in a clinical study. However a pronounced lowering of the cerebral seizure threshold may occur when quinolones are given concurrently with theophylline, non-steroidal anti-inflammatory drugs, or other agents which lower the seizure threshold.

Levofloxacin concentrations were about 13% higher in the presence of fenbufen than when administered alone.



Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

#### *Probenecid and cimetidine*

Probenecid and cimetidine had a statistically significant effect on the elimination of levofloxacin. The renal clearance of levofloxacin was reduced by cimetidine (24%) and probenecid (34%). This is because both drugs are capable of blocking the renal tubular secretion of levofloxacin. However, at the tested doses in the study, the statistically significant kinetic differences are unlikely to be of clinical relevance.

Caution should be exercised when levofloxacin is coadministered with drugs that affect the tubular renal secretion such as probenecid and cimetidine, especially in renally impaired patients.

#### *Other relevant information*

Clinical pharmacology studies have shown that the pharmacokinetics of levofloxacin were not affected to any clinically relevant extent when levofloxacin was administered together with the following

drugs: calcium carbonate, digoxin, glibenclamide, ranitidine.

#### Effect of Levo-Denk 500 on other medicinal products

##### *Ciclosporin*

The half-life of ciclosporin was increased by 33% when coadministered with levofloxacin.

##### *Vitamin K antagonists*

Increased coagulation tests (PT/INR) and/or bleeding, which may be severe, have been reported in patients treated with levofloxacin in combination with a vitamin K antagonist (e.g. warfarin). Coagulation tests, therefore, should be monitored in patients treated with vitamin K antagonists (see section 4.4).

##### *Drugs known to prolong QT interval*

Levofloxacin, like other fluoroquinolones, should be used with caution in patients receiving drugs known to prolong the QT interval (e.g. Class IA and III antiarrhythmics, tricyclic antidepressants, macrolides, antipsychotics) (see section 4.4 QT interval prolongation).

#### *Other relevant information*

In a pharmacokinetic interaction study, levofloxacin did not affect the pharmacokinetics of theophylline (which is a probe substrate for CYP1A2), indicating that levofloxacin is not a CYP1A2 inhibitor.

#### Other forms of interactions

##### *Food*

There is no clinically relevant interaction with food. Levo-Denk 500 tablets may therefore be administered regardless of food intake.

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

#### 4.6 Fertility, pregnancy and lactation

##### *Pregnancy*

There are limited amount of data from the use of levofloxacin in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3). However in the absence of human data and due to that experimental data suggest a risk of damage by fluoroquinolones to the weight-bearing cartilage of the growing organism, levofloxacin must not be used in pregnant women (see sections 4.3 and 5.3).

##### *Breast-feeding*

Levo-Denk 500 is contraindicated in breast-feeding women. There is insufficient information on the excretion of levofloxacin in human milk; however other fluoroquinolones are excreted in breast milk. In the absence of human data and due to that experimental data suggest a risk of damage by fluoroquinolones to the weight-bearing cartilage of the growing organism, levofloxacin must not be used in breast-feeding women (see sections 4.3 and 5.3).

##### *Fertility*

Levofloxacin caused no impairment of fertility or reproductive performance in rats.

#### 4.7 Effects on ability to drive and use machines

Some undesirable effects (e.g. dizziness/vertigo, drowsiness, visual disturbances) may impair the patient's ability to concentrate and react, and therefore may constitute a risk in situations where these abilities are of special importance (e.g. driving a car or operating machinery).

#### 4.8 Undesirable effects

The information given below is based on data from clinical studies in more than 8300 patients and on extensive post marketing experience.

Frequencies are defined using the following convention: very common ( $\geq 1/10$ ), common ( $\geq 1/100$ ,  $< 1/10$ ), uncommon ( $\geq 1/1000$ ,  $< 1/100$ ), rare ( $\geq 1/10000$ ,  $< 1/1000$ ), very rare ( $< 1/10000$ ), not known (cannot be estimated from the available data).

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

System organ class	Common ( $\geq 1/100$ to $< 1/10$ )	Uncommon ( $\geq 1/1,000$ to $< 1/100$ )	Rare ( $\geq 1/10,000$ to $< 1/1,000$ )	Not known (cannot be estimated from available data)
Infections and infestations		Fungal infection including Candida infection Pathogen resistance		
Blood and lymphatic system disorders		Leukopenia Eosinophilia	Thrombocytopenia Neutropenia	Pancytopenia Agranulocytosis Haemolytic anaemia

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

Immune system disorders			Angioedema Hypersensitivity (see section 4.4)	Anaphylactic shock <sup>a</sup> Anaphylactoid shock <sup>a</sup> (see section 4.4)
Metabolism and nutrition disorders		Anorexia	Hypoglycaemia particularly in diabetic patients (see section 4.4)	Hyperglycaemia Hypoglycaemic coma (see section 4.4)
Psychiatric disorders	Insomnia	Anxiety Confusional state Nervousness	Psychotic reactions (with e.g. hallucination, paranoia) Depression Agitation Abnormal dreams Nightmares	Psychotic disorders with self-endangering behaviour including suicidal ideation or suicide attempt (see section 4.4)
Nervous system disorders	Headache Dizziness	Somnolence Tremor Dysgeusia	Convulsion (see sections 4.3 and 4.4) Paraesthesia	Peripheral sensory neuropathy (see section 4.4) Peripheral sensory motor neuropathy (see section 4.4) Parosmia including anosmia Dyskinesia Extrapyramidal disorder Ageusia Syncope Benign intracranial hypertension
Eye disorders			Visual disturbances such as blurred vision (see section 4.4)	Transient vision loss (see section 4.4)
Ear and Labyrinth disorders		Vertigo	Tinnitus	Hearing loss Hearing impaired
Cardiac disorders			Tachycardia, Palpitation	Ventricular tachycardia, which may result in cardiac arrest Ventricular arrhythmia and torsade de pointes (reported predominantly in patients with risk factors of QT prolongation), electrocardiogram QT prolonged (see sections 4.4 and 4.9)
Vascular disorders	<u>Applies to iv form only:</u> Phlebitis		Hypotension	

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

Respiratory, thoracic and mediastinal disorders		Dyspnoea		Bronchospasm Pneumonitis allergic
Gastro-intestinal disorders	Diarrhoea Vomiting Nausea	Abdominal pain Dyspepsia Flatulence Constipation		Diarrhoea – haemorrhagic which in very rare cases may be indicative of enterocolitis, including pseudomembranous colitis (see section 4.4) Pancreatitis
Hepatobiliary disorders	Hepatic enzyme increased (ALT/AST, alkaline phosphatase, GGT)	Blood bilirubin increased		Jaundice and severe liver injury, including cases with fatal acute liver failure, primarily in patients with severe underlying diseases (see section 4.4) Hepatitis
Skin and subcutaneous tissue disorders <sup>b</sup>		Rash Pruritus Urticaria Hyperhidrosis		Toxic epidermal necrolysis Stevens-Johnson syndrome Erythema multiforme Photosensitivity reaction (see section 4.4) Leukocytoclastic vasculitis Stomatitis
Musculoskeletal and connective tissue disorders		Arthralgia Myalgia	Tendon disorders (see sections 4.3 and 4.4) including tendinitis (e.g. Achilles tendon) Muscular weakness which may be of special importance in patients with myasthenia gravis (see section 4.4 )	Rhabdomyolysis Tendon rupture (e.g. Achilles tendon) (see sections 4.3 and 4.4) Ligament rupture Muscle rupture Arthritis
Renal and Urinary disorders		Blood creatinine increased	Renal failure acute (e.g. due to interstitial nephritis)	
General disorders and administration site conditions	<u>Applies to iv form only:</u> Infusion site reaction (pain, reddening)	Asthenia	Pyrexia	Pain (including pain in back, chest, and extremities)

<sup>a</sup> Anaphylactic and anaphylactoid reactions may sometimes occur even after the first dose

<sup>b</sup> Mucocutaneous reactions may sometimes occur even after the first dose

Other undesirable effects which have been associated with fluoroquinolone administration include:

- attacks of porphyria in patients with porphyria

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

#### **4.9 Overdose**

According to toxicity studies in animals or clinical pharmacology studies performed with supra-therapeutic doses, the most important signs to be expected following acute overdose of Levo-Denk 500 tablets are central nervous system symptoms such as confusion, dizziness, impairment of consciousness, and convulsive seizures, increases in QT interval as well as gastro-intestinal reactions such as nausea and mucosal erosions.

CNS effects including confusional state, convulsion, hallucination, and tremor have been observed in post marketing experience.

In the event of overdose, symptomatic treatment should be implemented. ECG monitoring should be undertaken, because of the possibility of QT interval prolongation. Antacids may be used for protection of gastric mucosa. Haemodialysis, including peritoneal dialysis and CAPD, are not effective in removing levofloxacin from the body. No specific antidote exists.

### **5. PHARMACOLOGICAL PROPERTIES**

#### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: quinolone antibacterials, fluoroquinolones, ATC code: J01MA12  
Levofloxacin is a synthetic antibacterial agent of the fluoroquinolone class and is the S (-) enantiomer of the racemic active substance ofloxacin.

##### *Mechanism of action*

As a fluoroquinolone antibacterial agent, levofloxacin acts on the DNA-DNA-gyrase complex and topoisomerase IV.

##### *PK/PD relationship*

The degree of the bactericidal activity of levofloxacin depends on the ratio of the maximum concentration in serum (C<sub>max</sub>) or the area under the curve (AUC) and the minimal inhibitory concentration (MIC).

##### *Mechanism of resistance*

Resistance to levofloxacin is acquired through a stepwise process by target site mutations in both type II topoisomerases, DNA gyrase and topoisomerase IV. Other resistance mechanisms such as permeation barriers (common in *Pseudomonas aeruginosa*) and efflux mechanisms may also affect susceptibility to levofloxacin.

Cross-resistance between levofloxacin and other fluoroquinolones is observed. Due to the mechanism of action, there is generally no cross-resistance between levofloxacin and other classes of antibacterial agents.

##### *Breakpoints*

The EUCAST recommended MIC breakpoints for levofloxacin, separating susceptible from intermediately susceptible organisms and intermediately susceptible from resistant organisms are presented in the below table for MIC testing (mg/l).

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

EUCAST clinical MIC breakpoints for levofloxacin (version 2.0, 2012-01-01):

Pathogen	Susceptible	Resistant
Enterobacteriaceae	≤1 mg/l	>2 mg/l
<i>Pseudomonas spp.</i>	≤1 mg/l	>2 mg/l
<i>Acinetobacter spp.</i>	≤1 mg/l	>2 mg/l
<i>Staphylococcus spp.</i>	≤1 mg/l	>2 mg/l
<i>S. pneumoniae</i> <sup>1</sup>	≤2 mg/l	>2 mg/l
<i>Streptococcus A, B, C, G</i>	≤1 mg/l	>2 mg/l
<i>H. influenzae</i> <sup>2,3</sup>	≤1 mg/l	>1 mg/l
<i>M. catarrhalis</i> <sup>3</sup>	≤1 mg/l	>1 mg/l
Non-species related breakpoints <sup>4</sup>	≤1 mg/l	>2 mg/l

1. The breakpoints for levofloxacin relate to high dose therapy.  
2. Low-level fluoroquinolone resistance (ciprofloxacin MICs of 0.12-0.5 mg/l) may occur but there is no evidence that this resistance is of clinical importance in respiratory tract infections with *H. influenzae*.  
3. Strains with MIC values above the susceptible breakpoint are very rare or not yet reported. The identification and antimicrobial susceptibility tests on any such isolate must be repeated and if the result is confirmed the isolate must be sent to a reference laboratory. Until there is evidence regarding clinical response for confirmed isolates with MIC above the current resistant breakpoint they should be reported resistant.  
4. Breakpoints apply to an oral dose of 500 mg x 1 to 500 mg x 2 and an intravenous dose of 500 mg x 1 to 500 mg x 2.

The prevalence of resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert advice should be sought when the local prevalence of resistance is such that the utility of the agent in at least some types of infections is questionable.

### **Commonly susceptible species**

#### **Aerobic Gram-positive bacteria**

*Bacillus anthracis*  
*Staphylococcus aureus* methicillin-susceptible  
*Staphylococcus saprophyticus*  
*Streptococci, group C and G*  
*Streptococcus agalactiae*  
*Streptococcus pneumoniae*  
*Streptococcus pyogenes*

#### **Aerobic Gram-negative bacteria**

*Eikenella corrodens*  
*Haemophilus influenzae*  
*Haemophilus para-influenzae*  
*Klebsiella oxytoca*  
*Moraxella catarrhalis*  
*Pasteurella multocida*  
*Proteus vulgaris*  
*Providencia rettgeri*

#### **Anaerobic bacteria**

*Peptostreptococcus*

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

### **Other**

*Chlamydophila pneumoniae*  
*Chlamydophila psittaci*  
*Chlamydia trachomatis*  
*Legionella pneumophila*  
*Mycoplasma pneumoniae*  
*Mycoplasma hominis*  
*Ureaplasma urealyticum*

### **Species for which acquired resistance may be a problem**

#### **Aerobic Gram-positive bacteria**

*Enterococcus faecalis*  
*Staphylococcus aureus* methicillin-resistant#  
Coagulase negative *Staphylococcus spp*

#### **Aerobic Gram- negative bacteria**

*Acinetobacter baumannii*  
*Citrobacter freundii*  
*Enterobacter aerogenes*  
*Enterobacter cloacae*  
*Escherichia coli* *Klebsiella pneumoniae* *Morganella morganii* *Proteus mirabilis*  
*Providencia stuartii*  
*Pseudomonas aeruginosa*  
*Serratia marcescens*

#### **Anaerobic bacteria**

*Bacteroides fragilis*

### **Inherently Resistant Strains**

#### **Aerobic Gram-positive bacteria**

*Enterococcus faecium*

# Methicillin-resistant *S. aureus* are very likely to possess co-resistance to fluoroquinolones, including levofloxacin.

## **5.2 Pharmacokinetic properties**

### **Absorption**

Orally administered levofloxacin is rapidly and almost completely absorbed with peak plasma concentrations being obtained within 1 - 2 h. The absolute bioavailability is 99 - 100%.

Food has little effect on the absorption of levofloxacin.

Steady state conditions are reached within 48 hours following a 500 mg once or twice daily dosage regimen.

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

### Distribution

Approximately 30 - 40% of levofloxacin is bound to serum protein.

The mean volume of distribution of levofloxacin is approximately 100 l after single and repeated 500 mg doses, indicating widespread distribution into body tissues.

### Penetration into tissues and body fluids

Levofloxacin has been shown to penetrate into bronchial mucosa, epithelial lining fluid, alveolar macrophages, lung tissue, skin (blister fluid), prostatic tissue and urine. However, levofloxacin has poor penetration into cerebro-spinal fluid.

### Biotransformation

Levofloxacin is metabolised to a very small extent, the metabolites being desmethyl-levofloxacin and levofloxacin N-oxide. These metabolites account for <5% of the dose and are excreted in urine. Levofloxacin is stereochemically stable and does not undergo chiral inversion.

### Elimination

Following oral and intravenous administration of levofloxacin, it is eliminated relatively slowly from the plasma ( $t_{1/2}$ : 6 - 8 h). Excretion is primarily by the renal route (>85% of the administered dose).

The mean apparent total body clearance of levofloxacin following a 500 mg single dose was 175 +/-29.2 ml/min.

There are no major differences in the pharmacokinetics of levofloxacin following intravenous and oral administration, suggesting that the oral and intravenous routes are interchangeable.

### Linearity

Levofloxacin obeys linear pharmacokinetics over a range of 50 to 1000 mg.

### Special populations

#### *Subjects with renal insufficiency*

The pharmacokinetics of levofloxacin are affected by renal impairment. With decreasing renal function renal elimination and clearance are decreased, and elimination half-lives increased as shown in the table below:

Pharmacokinetics in renal insufficiency following single oral 500 mg dose

Cl <sub>cr</sub> [ml/min]	<20	20 - 49	50 - 80
Cl <sub>R</sub> [ml/min]	13	26	57
t <sub>1/2</sub> [h]	35	27	9

#### *Elderly subjects*

There are no significant differences in levofloxacin pharmacokinetics between young and elderly subjects, except those associated with differences in creatinine clearance.



Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

### *Gender differences*

Separate analysis for male and female subjects showed small to marginal gender differences in levofloxacin pharmacokinetics. There is no evidence that these gender differences are of clinical relevance.

### **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on conventional studies of single dose toxicity, repeated dose toxicity, carcinogenic potential and toxicity to reproduction and development.

Levofloxacin caused no impairment of fertility or reproductive performance in rats and its only effect on fetuses was delayed maturation as a result of maternal toxicity.

Levofloxacin did not induce gene mutations in bacterial or mammalian cells but did induce chromosome aberrations in Chinese hamster lung cells *in vitro*. These effects can be attributed to inhibition of topoisomerase II. *In vivo* tests (micronucleus, sister chromatid exchange, unscheduled DNA synthesis, dominant lethal tests) did not show any genotoxic potential.

Studies in the mouse showed levofloxacin to have phototoxic activity only at very high doses.

Levofloxacin did not show any genotoxic potential in a photomutagenicity assay, and it reduced tumour development in a photocarcinogenicity study.

In common with other fluoroquinolones, levofloxacin showed effects on cartilage (blistering and cavities) in rats and dogs. These findings were more marked in young animals.

## **6. Pharmaceutical particulars**

### **6.1 List of excipients**

Tablet core: Microcrystalline cellulose, powdered cellulose, pregelatinised starch (maize), maize starch, crospovidone, povidone K25, sodium stearyl fumarate

Tablet coating: Lactose monohydrate, hypromellose, titanium dioxide (E 171), macrogol 4000, ferric oxides and ferric hydroxides (E 172)

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years

### **6.4 Special precautions for storage**

Store in a dry place below 30 °C.

Do not use after the expiry date.

Keep out of the reach and sight of children.

Trade name:	Levo-Denk 500
Generic name:	Levofloxacin 500 mg, film-coated tablet
Applicant:	DENK PHARMA GmbH & Co. KG

### **6.5 Nature and contents of container**

PVC/PVDC aluminium blister strips.  
Pack size: 5 and 10 film-coated tablets  
Free medical samples: 5 and 10 film-coated tablets

### **6.6 Special precautions for disposal**

A score line allows adaptation of the dose in patients with impaired renal function.  
As for all medicines, any unused product or waste material should be disposed of in accordance with local environmental requirements.

### **7. Marketing authorisation holder**

DENK PHARMA GmbH & Co. KG  
Prinzregentenstr. 79  
81675 München  
Germany

### **8. Marketing authorisation number in Germany**

70183.00.00

### **9. Date of first authorisation in Germany**

17.12.2008

### **10. Date of revision of the text**

02/2014

### **11. Prescription status**

Prescription only medicine