

**SUMMARY OF PRODUCT CHARACTERISTICS (SPC)**

## 1. NAME OF THE MEDICINAL PRODUCT

**ONCODAC** (Dacarbazine 200mg lyophilized powder for injection)

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial contains 200 mg dacarbazine.

*Excipient(s) with known effect:*

Each ml contains 100 mg mannitol.

For the full list of excipients, see section 6.1

## 3. PHARMACEUTICAL FORM

Lyophilized powder for injection

White to pale yellow lyophilized cake

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Dacarbazine is indicated for the treatment of patients with metastasised malignant melanoma.

Further indications for dacarbazine as part of a combination chemotherapy are:

- advanced Hodgkin's disease,
- advanced adult soft tissue sarcomas (except mesothelioma, Kaposi sarcoma).

### 4.2 Posology and method of administration

#### Posology

The use of dacarbazine should be confined to physicians experienced in oncology or haematology.

Dacarbazine is sensitive to light exposure. All reconstituted solutions should be suitably protected from light also during administration (light-resistant infusion set).

Care should be taken when administering the injection to avoid extravasation into tissues since this will cause local pain and tissue damage. If extravasation occurs, the injection should be discontinued immediately and any remaining portion of the dose should be introduced into another vein.

The following regimes may be used. For further details see current scientific literature.

#### *Malignant melanoma*

Dacarbazine can be administered as single agent in doses of 200 to 250 mg/m<sup>2</sup> body surface area/day as an i.v. injection for 5 days every 3 weeks.

As an alternative to an intravenous bolus injection dacarbazine can be administered as a short-term infusion (over 15 – 30 minutes).

It is also possible to give 850 mg/m<sup>2</sup> body surface area on day 1 and then once every 3 weeks as intravenous infusion.

### *Hodgkin's disease*

Dacarbazine is administered in a daily dose of 375 mg/m<sup>2</sup> body surface area i.v. every 15 days in combination with doxorubicin, bleomycin and vinblastine (ABVD regimen).

### *Adult soft-tissue sarcoma*

For adult soft tissue sarcomas dacarbazine is given in daily doses of 250 mg/m<sup>2</sup> body surface area i.v. (days 1 - 5) in combination with doxorubicin every 3 weeks (ADIC regimen).

During dacarbazine treatment frequent monitoring of blood counts should be conducted as well as monitoring of hepatic and renal function. Since severe gastrointestinal reactions frequently occur, antiemetic and supportive measures are advisable.

Because severe gastrointestinal and haematological disturbances can occur an extremely careful benefit-risk analysis has to be made before every course of therapy with dacarbazine.

### *Duration of therapy*

The treating physician should individually decide about the duration of therapy taking into account the type and stage of the underlying disease, the combination therapy administered and the response to and adverse effects of dacarbazine. In advanced Hodgkin's disease, a usual recommendation is to administer 6 cycles of ABVD combination therapy. In metastasised malignant melanoma and in advanced tissue sarcoma, the duration of treatment depends on the efficacy and tolerability in the individual patient.

### *Patients with kidney/liver insufficiency*

If there is mild to moderate renal or hepatic insufficiency alone, a dose reduction is not usually required. In patients with combined renal and hepatic impairment elimination of dacarbazine is prolonged. However, no validated recommendations on dose reductions can be given currently.

### *Elderly patients*

As limited experience in elderly patients is available no special instructions for the use in elderly patients can be given.

### *Paediatric population*

The safety and efficacy of dacarbazine in children/adolescents aged < 15 years have not yet been established. No special recommendations for the use of dacarbazine in the paediatric age group can be given until further data become available.

## Method of administration

### *Rate of administration*

Doses up to 200 mg/m<sup>2</sup> may be given as a slow intravenous injection. Larger doses (ranging from 200 to 850 mg/m<sup>2</sup>) should be administered as an i.v. infusion over 15 – 30 minutes.

It is recommended to test the patency of the vein first with a 5- to 10-ml flush of 0.9 % sodium chloride or 5 % glucose infusion solution. The same solutions should be used after infusion to flush any remaining medicinal product from the tubing.

After reconstitution with water for injections without further dilution with 0.9 % sodium chloride or 5 % glucose infusion solution, dacarbazine 100 mg and 200 mg preparations are hypo-osmolar (ca. 100 mOsmol/kg) and should therefore be given by slow intravenous injection e.g. over 1 minute rather than rapid intravenous bolus over a few seconds.

*Precautions to be taken before handling or administering the medicinal product*

### **4.3 Contraindications**

- Hypersensitivity to the active substance or to any of the excipients
- pregnancy or breastfeeding
- leukopenia and/or thrombocytopenia,
- severe liver or kidney diseases.

### **4.4 Special warnings and precautions for use**

It is recommended that dacarbazine should only be administered under the supervision of a physician specialised in oncology who has the facilities for regular monitoring of clinical, biochemical and haematological effects, during and after therapy.

If symptoms of a liver or kidney functional disorder or symptoms of a hypersensitivity reaction are observed immediate cessation of therapy is required. If veno-occlusive disease of the liver occurs, further therapy with dacarbazine is contraindicated.

Note: The responsible physician should be aware of a rarely observed severe complication during therapy resulting from liver necrosis due to occlusion of intrahepatic veins. Therefore frequent monitoring of liver size, function and blood counts (especially eosinophils) is required. In single cases of suspected veno-occlusive disease early therapy with high-dose corticosteroids (for example hydrocortisone 300 mg/day) with or without fibrinolytic agents like heparin or tissue plasminogen activator was successful.

Long-term therapy can cause cumulative bone marrow toxicity. The possible bone marrow depression requires careful monitoring of white blood cells, red blood cells and platelet levels. Haemopoietic toxicity may warrant temporary suspension or cessation of therapy.

Extravasation of the medicinal product during i.v. administration may result in tissue damage and severe pain.

Concomitant use with phenytoin should be avoided because reduced absorption of phenytoin from the gastrointestinal tract may predispose the patient to convulsions.

Dacarbazine is a moderate immunosuppressive agent. Administration of live vaccines to patients who are immunocompromised as a result of treatment with chemotherapeutics such as dacarbazine can cause serious and potentially fatal infections. Immunisation with live vaccines should therefore be avoided during dacarbazine therapy. It is generally advised to use live virus vaccines with caution after stopping chemotherapy and to take the patient's immune status into account, depending also on the disease and other therapies. Vaccination with live vaccines should be administered no sooner than 3 months after the completion of chemotherapy. Inactivated vaccines can be used if available.

Concomitant use of fotemustine can cause acute pulmonary toxicity (adult respiratory distress syndrome), which may lead to a fatal outcome. Fotemustine and dacarbazine should not be used concomitantly.

Hepatotoxic medicinal products and alcohol should be avoided during chemotherapy.

#### Contraceptive measures

Men are advised to take contraceptive measures during and for 6 months after cessation of therapy.

#### Paediatric population

Dacarbazine is not recommended for use in the paediatric age group until further data become available.

### **4.5 Interaction with other medicinal products and other forms of Interaction**

In case of previous or concomitant treatment having adverse effects on the bone marrow (particularly cytostatic agents, irradiation) myelotoxic interactions are possible.

Studies to investigate the presence of phenotypic metabolism have not been undertaken but hydroxylation of the parent compound to metabolites with anti-tumour activity has been identified.

Dacarbazine is metabolised by cytochrome P450 (CYP1A1, CYP1A2, and CYP2E1). This has to be taken into account if other medicinal products are co-administered which are metabolised by the same hepatic enzymes.

Dacarbazine can enhance the effects of methoxypsoralen because of photosensitization.

Immunisation with live vaccines should be avoided during therapy with dacarbazine due to the risk of serious and potentially fatal infections. It is advised to use live virus vaccines with caution after stopping chemotherapy and vaccinate not sooner than 3 months after the last dose of chemotherapy. It is recommended to use an inactivated vaccine if available.

Risk of thrombosis is increased in malignant diseases; therefore, use of concomitant anticoagulation is common. If the patient is to receive oral anticoagulants, the frequency of INR monitoring must be increased due to large inter individual variability in coagulation and due to possible interaction between anticoagulants and cytostatics.

Concomitant use with phenytoin may cause reduced absorption of phenytoin from the gastrointestinal tract and may predispose the patient to convulsions.

Concomitant use of cyclosporine (and in some cases tacrolimus) must be considered carefully because these agents may cause excessive immunosuppression and lymphoproliferation.

Concomitant use of fotemustine can cause acute pulmonary toxicity (adult respiratory distress syndrome). Fotemustine and dacarbazine should not be used concomitantly.

#### 4.6 Pregnancy and lactation

##### Pregnancy

Dacarbazine has been shown to be mutagenic, teratogenic and carcinogenic in animals. It must be assumed that an increased risk for teratogenic effects exists in humans. Therefore Dacarbazine is contraindicated during pregnancy.

Women of child-bearing potential have to use effective contraception during treatment.

##### Breast-feeding

Dacarbazine is contraindicated during breast-feeding

#### 4.7 Undesirable effects

##### Frequencies

Very common (> 1/10)

Common (> 1/100 to < 1/10)

Uncommon (> 1/1000 to < 1/100)

Rare (> 1/10000 to < 1/1000)

Very rare (< 1/10000)

Not known (cannot be estimated from the available data)

The most commonly reported ADRs are gastrointestinal disorders (anorexia, nausea and vomiting) and blood and lymphatic system disorders such as anaemia, leukopenia and thrombocytopenia. The latter are dose-dependent and delayed, with the nadirs often only occurring after 3 to 4 weeks.

Infections and infestations	<u>Uncommon</u> Infections
Blood and lymphatic system disorders	<u>Common</u> Anaemia, leukopenia, thrombocytopenia <u>Rare</u> <u>Pancytopenia, agranulocytosis</u>
Immune system disorders	<u>Rare</u> Anaphylactic reactions
Nervous system disorders	<u>Rare</u> Headaches, impaired vision, confusion, lethargy,

	convulsions, facial paraesthesia
Vascular disorders	<u>Rare</u> <u>Facial flushing</u>
Gastrointestinal disorders	<u>Common</u> Anorexia, nausea, vomiting <u>Rare</u> Diarrhoea
Hepatobiliary disorders	<u>Rare</u> Hepatic necrosis due to veno-occlusive disease (VOD) of the liver, Budd-Chiari syndrome (with potentially fatal outcome)
Renal and urinary disorders	<u>Rare</u> Impaired renal function
Skin and subcutaneous tissue disorders	<u>Uncommon</u> Alopecia, hyperpigmentation, photosensitivity <u>Rare</u> Erythema, maculopapular exanthema, urticaria
General disorders and administration site conditions	<u>Uncommon</u> Flu-like symptoms <u>Rare</u> Application site irritation
Investigations	<u>Rare</u> Hepatic enzymes increased (e.g. alkaline phosphatase, ASAT, ALAT), blood lactate dehydrogenase (LDH) increased, blood creatinine increased, blood urea increased

### Description of selected adverse reactions

Changes in blood counts often observed (anaemia, leukopenia, thrombocytopenia) are dose-dependent and delayed, with the nadirs often only occurring after 3 to 4 weeks.

Flu-like symptoms with exhaustion, chills, fever and muscular pain are occasionally observed during or often only days after dacarbazine administration. These disturbances may recur with the next infusion.

Rarely liver necrosis due to occlusion of intrahepatic veins (veno-occlusive disease of the liver) has been observed after administration of dacarbazine in monotherapy or in combined treatment modalities. In general the syndrome occurred during the second cycle of therapy. Symptoms included fever, eosinophilia, abdominal pain, enlarged liver, jaundice and shock which worsened rapidly over a few hours or days. As fatal outcome has been described special care has to be taken.

Application site irritations and some of the systemic adverse reactions are thought to result from formation of photodegradation products.

Facial paraesthesia and flushing may occur shortly after injection.

Allergic reactions of the skin in the form of erythema, maculopapular exanthema or urticaria are observed rarely.

Inadvertent paravenous injection is expected to cause local pain and necrosis.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions.

### **4.8 Overdose**

The primary anticipated complications of overdose are severe bone marrow suppression, eventually bone marrow aplasia which may be delayed by up to two weeks. Time to occurrence of nadirs of leucocytes and thrombocytes can be 4 weeks. Even if overdose is only suspected, long-term careful haematologic monitoring is essential.

There is no known antidote for dacarbazine overdose. Therefore, special care has to be taken to avoid overdose of this medicinal product.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Alkylating agents, ATC code: L01AX04.

Dacarbazine is a cytostatic agent. The antineoplastic effect is due to an inhibition of cell growth which is independent of the cell cycle and due to an inhibition of DNA synthesis. An alkylating effect has also been shown and other cytostatic mechanisms may also be influenced by dacarbazine.

Dacarbazine is considered not to show an antineoplastic effect by itself. However by microsomal N-demethylation it is quickly converted to 5-amino-imidazole-4-carboxamide and a methyl cation, which is responsible for the alkylating effect of the medicinal product.

### **5.2 Pharmacokinetic properties**

#### Distribution

After intravenous administration dacarbazine is quickly distributed into tissue. Plasma protein binding is 5 %. Kinetics in plasma are biphasic; the initial (distribution) half-life is only 20 minutes, terminal half-life is 0.5 – 3.5 hours.

#### Biotransformation

Dacarbazine is inactive until metabolised in the liver by cytochromes P450 to form the reactive N demethylated species HMMTIC and MTIC. This is catalysed by CYP1A1, CYP1A2, and CYP2E1.



MTIC is further metabolised to 5-aminoimidazole-4-carboxamide (AIC).

### Elimination

Dacarbazine is metabolised mainly in the liver by both hydroxylation and demethylation, approx. 20 – 50 % of the medicinal product is excreted unmodified by the kidney via renal tubular secretion.

### **5.3 Preclinical safety data**

Because of its pharmacodynamic properties dacarbazine shows mutagenic, carcinogenic and teratogenic effects which are detectable in experimental test systems.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of Excipients**

Citric Acid

Mannitol

Sodium Hydroxide

Water for Injection

### **6.2 Incompatibilities**

Dacarbazine solution is chemically incompatible with heparin, hydrocortisone, L-cysteine and sodium hydrogen carbonate.

### **6.3 Shelf life**

24 Months

### **6.4 Storage conditions**

Store between 2°C to 8°C, protect from light. DO NOT FREEZE.

### **6.5 Nature and contents of container**

20 ml Amber tubular glass vial with 20 mm rubber plug sealed with 20 mm Flip of Aluminium Seal is placed in a carton along with pack insert.

### **6.6 Special precautions for disposal and other handling**

#### Recommendations for safe handling

Dacarbazine is an antineoplastic agent and should be handled according to standard procedures for cytostatics that have mutagenic, carcinogenic and teratogenic effects. Before commencing, local cytotoxic guidelines should be referred to.

Dacarbazine should only be opened by trained staff and as with all cytotoxic agents; precautions should be taken to avoid exposing staff. Handling of cytotoxic medicinal products should be generally avoided during pregnancy. Preparation of solution for administration should be carried out in a designated handling area and working over a washable tray or disposable plastic-backed absorbent paper.

Suitable eye protection, disposable gloves, face mask and disposable apron should be worn. Syringes and infusion sets should be assembled carefully to avoid leakage (use of Luer lock fittings is recommended).

On completion, any exposed surface should be thoroughly cleaned and hands and face washed.

In the event of spillage, operators should put on gloves, face masks, eye-protection and disposable apron and mop up the spilled material with an absorbent material tamped in the area for that purpose. The area should then be cleaned and all contaminated material transferred to a cytotoxic spillage bag or bin or sealed for incineration.

#### Preparation for intravenous administration

Dacarbazine-solutions are prepared immediately before use.

Dacarbazine is sensitive to light exposure. During administration, the infusion container and administration set should be protected from exposure to daylight, e.g. by using light-resistant PVC-infusion sets. Normal infusion sets should be wrapped up in e.g. UV-resistant foils.

#### **Preparation of Dacarbazine 200 mg:**

Aseptically transfer 20 ml of water for injections into the vial and shake until a solution is obtained. This freshly prepared solution, containing 10 mg/ml of dacarbazine, (density of the solution:  $\rho = 1.007$  g/ml) is administered as a slow injection.

For preparation of Dacarbazine 200 mg for i.v. infusion the freshly prepared solution is further diluted with 200 – 300 ml 0.9 % sodium chloride or 5 % glucose infusion solution. This solution is given as a short term infusion over a period between 15 – 30 minutes.

Diluted solution should be visually inspected and only clear solutions practically free from particles should be used. Do not use solution if particles are present.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

#### **7. Marketing Authorization Holder**

M/s VHB Medi Sciences Ltd.

Plot No- 20-22 & 49-51, IIE, Sector-5

Pantnagar, Udham Singh Nagar

Uttarakhand, INDIA

Tel. No.: +915944-666200

**Manufacturer**

M/s VHB Medi Sciences Ltd.

Plot No-50 AB, Govt Industrial Estate,

Charkop, Kandivali(W)

Mumbai 400 067

INDIA

Tel. No.: +91 22 4163 9000

Fax No.: +91 22 4163 9100

**8. MARKETING AUTHORISATION NUMBER**

05997/07285/REN/2020

**9. DATE OF FIRST REGISTRATION/RENEWAL OF THE REGISTRATION**

24/05/2021

**10. DATE OF REVISION OF THE TEXT**

18/07/2023