

## **SUMMARY OF PRODUCT CHARACTERISTICS**

## **1. Name of the medicinal product**

**BRUZOLIN NASAL DROPS (Xylometazoline Nasal Drops BP 0.1% w/v)**

## **2. Qualitative and quantitative composition**

Composition:

Xylometazoline Hydrochloride BP .....0.1% w/v

Benzalkonium Chloride Solution BP .....0.02% w/v

Purified water BP .....q.s

## **3. Pharmaceutical form**

Nasal Drops

Clear colorless solution filled in white LDPE Bottle.

## **4. Clinical particulars**

### **4.1 Therapeutic indications**

For the relief of nasal congestion associated with the common cold, perennial and allergic rhinitis (including hay fever), sinusitis.

### **4.2 Posology and method of administration**

Adults and elderly (all indications): 2 or 3 drops in each nostril up to 3 times daily. Do not exceed 3 applications daily into each nostril.

The drops are suitable for children over 12 years of age. The recommended dose should not be exceeded, especially in children and the elderly.

Route of administration: Nasal use.

### **4.3 Contraindications**

Known hypersensitivity to Bruzolin

Patients with trans-sphenoidal hypophysectomy or surgery exposing the dura mater

Narrow-angle glaucoma

Rhinitis sicca or atrophic rhinitis.

Bruzolin 0.1% is contraindicated in children aged less than 12 years

People with phaeochromocytoma or prostatic hypertrophy or receiving monoamine oxidase inhibitors (MAOI) treatment or who have received them in the last two weeks.

#### **4.4 Special warnings and precautions for use**

Patients are advised not to take decongestants for more than seven consecutive days. Prolonged or excessive use may cause rebound congestion and/or atrophy of the nasal mucosa.

Bruzolin, like other preparations belonging to the same class of active substances, should be used only with caution in patients showing a strong reaction to sympathomimetic agents as evidenced by signs of insomnia, dizziness, tremor, cardiac arrhythmias or elevated blood pressure.

Caution is recommended in patients with hypertension, cardiovascular disease, hyperthyroidism or diabetes mellitus.

Patients with long QT syndrome treated with xylometazoline may be at increased risk of serious ventricular arrhythmias.

Keep medicines out of the sight and reach of children

#### **Information concerning excipients**

Bruzolin contains benzalkonium chloride. This may cause irritation of the nasal mucosa.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

The concomitant use of xylometazoline with monoamine oxidase (MAO) inhibitors or tri- and tetra-cyclic antidepressants, may cause an increase in blood pressure due to the cardiovascular effects of these substances (*see Contraindications*).

#### **4.6 Fertility, Pregnancy and lactation**

No foetal toxicity or fertility studies have been carried out in animals. In view of its potential systemic vasoconstrictor effect, it is advisable to take the precaution of not using Bruzolin during pregnancy.

No evidence of any adverse effect on the breast-fed infant. However, it is not known if xylometazoline is excreted in breast milk, therefore caution should be exercised and Bruzolin should be used only on the advice of a doctor whilst breastfeeding.

#### 4.7 Effects on ability to drive and use machines.

Bruzolin has no or negligible influence on the ability to drive and use machines.

#### 4.8 Undesirable effects

The adverse effects listed below are classified by system organ class and frequency according to the following convention: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ) or very rare ( $< 1/10,000$ ).

MeDRA SOC	Adverse reaction	Frequency
Immune System Disorders	Hypersensitivity reaction (angioedema, rash, pruritus)	Very rare
Nervous System Disorders	Headache	Common
Eye Disorders	Transient visual impairment	Very rare
Cardiac Disorders	Heart rate irregular Heart rate increased	Very rare Very rare
Respiratory, thoracic and mediastinal disorders	Nasal Dryness Nasal Discomfort Epistaxis	Common Common Uncommon
Gastrointestinal disorders	Nausea	Common
General disorders and administration site	Application site burning	Common

Other side effects include:

- A burning sensation in the nose and throat

#### 4.9 Overdose

##### Symptoms and signs

Excessive administration of topical xylometazoline hydrochloride or accidental ingestion may cause severe dizziness, perspiration, severely lowered body temperature, headache, bradycardia, hypertension, respiratory depression, coma and convulsions. Hypertension may be followed by hypotension. Small children are more sensitive to toxicity than adults.

##### Treatment

Appropriate supportive measures should be initiated in all individuals suspected of an overdose, and urgent symptomatic treatment under medical supervision is indicated when warranted. This would include observation of the individual for several hours.

## **5. Pharmacological properties**

### **5.1 Pharmacodynamic properties**

Bruzolin Nasal Drops is a sympathomimetic agent with marked alpha-adrenergic activity, and is intended for use in the nose. It constricts the nasal blood vessels, thereby decongesting the mucosa of the nose and neighbouring regions of the pharynx. This enables patients suffering from colds to breathe more easily through the nose. The effect of Bruzolin Nasal Drops begins within a few minutes and lasts for up to 10 hours. Bruzolin Nasal Drops is generally well tolerated and does not impair the function of ciliated epithelium.

In a double-blind, saline solution (Otrisal) controlled study in patients with common cold, the decongestant effect of Bruzolin was significantly superior ( $p < 0.0001$ ) to Otrisal saline solution based on rhinomanometry measurement at 1 hour after administration of the study drugs.

### **5.2 Pharmacokinetic properties**

Systemic absorption may occur following nasal application of xylometazoline hydrochloride solutions. It is not used systemically.

### **5.3 Preclinical safety data:**

There are no findings in the preclinical testing which are of relevance to the prescriber.

## **6. Pharmaceutical particulars**

### **6.1 List of excipients**

Benzalkonium chloride

Disodium edetate

Sodium chloride

Sodium citrate

Purified water

### **6.2 Incompatibilities**

Not applicable

**6.3 Shelf life**

24 Months

**6.4 Special precautions for storage**

Store at a temperature not exceeding 30°C. Keep out of reach of children.

**6.5 Nature and contents of container**

1 X 15ml LDPE Bottle packed in a carton along with package insert.

**6.6 Special precautions for disposal and other handling**

This medicinal product does not require any special storage conditions.

**7. Marketing authorisation holder**

Brawn Laboratories Limited.

Location (address): 13, N.I.T. Industrial Area,  
FARIDABAD-121001, (HARYANA)

Country: INDIA

Telephone: +91-129-4360113

E-Mail: [regulatory@brawnlabs.in](mailto:regulatory@brawnlabs.in)

Website: [www.brawnlabs.in](http://www.brawnlabs.in)

**8. Marketing authorisation number(s)**

06522/08105/REN/2021

**9. Date of first authorisation/renewal of the authorisation**

Sep 8, 2021

**10. Date of revision of the text**

18.07.2023