

1. NAME OF THE MEDICINAL PRODUCT

GONAL-f 450 IU/0.75 mL (33 micrograms/0.75 mL) solution for injection in pre-filled pen.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each mL of the solution contains 600 IU of follitropin alfa*, (equivalent to 44 micrograms).

Each pre-filled multidose pen delivers 450 IU (equivalent to 33 micrograms) in 0.75 mL.

* recombinant human follicle stimulating hormone (r-hFSH) produced in Chinese Hamster Ovary (CHO) cells by recombinant DNA technology.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for injection in pre-filled pen.

Clear colourless solution.

The pH of the solution is 6.7-7.3.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

In adult women

- Anovulation (including polycystic ovarian syndrome) in women who have been unresponsive to treatment with clomiphene citrate.
- Stimulation of multifollicular development in women undergoing superovulation for assisted reproductive technologies (ART) such as *in vitro* fertilisation (IVF), gamete intra-fallopian transfer and zygote intra-fallopian transfer.
- GONAL-f in association with a luteinising hormone (LH) preparation is recommended for the stimulation of follicular development in women with severe LH and FSH deficiency. In clinical trials these patients were defined by an endogenous serum LH level < 1.2 IU/L.

In adult men

- GONAL-f is indicated for the stimulation of spermatogenesis in men who have congenital or acquired hypogonadotropic hypogonadism with concomitant human Chorionic Gonadotropin (hCG) therapy.

4.2 Posology and method of administration

Treatment with GONAL-f should be initiated under the supervision of a physician experienced in the treatment of fertility disorders.

Patients must be provided with the correct number of pens for their treatment course and educated to use the proper injection techniques.

Posology

The dose recommendations given for GONAL-f are those in use for urinary FSH. Clinical assessment of GONAL-f indicates that its daily doses, regimens of administration, and treatment monitoring procedures should not be different from those currently used for urinary FSH-containing medicinal products. It is advised to adhere to the recommended starting doses indicated below.

Comparative clinical studies have shown that on average patients require a lower cumulative dose and shorter treatment duration with GONAL-f compared with urinary FSH. Therefore, it is considered appropriate to give a lower total dose of GONAL-f than generally used for urinary FSH, not only in order to optimise follicular development but also to minimise the risk of unwanted ovarian hyperstimulation. See section 5.1.

Bioequivalence has been demonstrated between equivalent doses of the monodose presentation and the multidose presentation of GONAL-f.

Women with anovulation (including polycystic ovarian syndrome)

GONAL-f may be given as a course of daily injections. In menstruating women treatment should commence within the first 7 days of the menstrual cycle.

A commonly used regimen commences at 75-150 IU FSH daily and is increased preferably by 37.5 or 75 IU at 7 or preferably 14 day intervals if necessary, to obtain an adequate, but not excessive, response. Treatment should be tailored to the individual patient's response as assessed by measuring follicle size by ultrasound and/or oestrogen secretion. The maximal daily dose is usually not higher than 225 IU FSH. If a patient fails to respond adequately after 4 weeks of treatment, that cycle should be abandoned and the patient should undergo further evaluation after which she may recommence treatment at a higher starting dose than in the abandoned cycle.

When an optimal response is obtained, a single injection of 250 micrograms recombinant human choriogonadotropin alfa (r-hCG) or 5,000 IU, up to 10,000 IU hCG should be administered 24-48 hours after the last GONAL-f injection. The patient is recommended to have coitus on the day of, and the day following, hCG administration. Alternatively intrauterine insemination (IUI) may be performed.

If an excessive response is obtained, treatment should be stopped and hCG withheld (see section 4.4). Treatment should recommence in the next cycle at a dose lower than that of the previous cycle.

Women undergoing ovarian stimulation for multiple follicular development prior to in vitro fertilisation or other assisted reproductive technologies.

A commonly used regimen for superovulation involves the administration of 150-225 IU of GONAL-f daily, commencing on days 2 or 3 of the cycle. Treatment is continued until adequate follicular development has been achieved (as assessed by monitoring of serum oestrogen concentrations and/or ultrasound examination), with the dose adjusted according to the patient's response, to usually not higher than 450 IU daily. In general adequate follicular development is achieved on average by the tenth day of treatment (range 5 to 20 days).

A single injection of 250 micrograms r-hCG or 5,000 IU up to 10,000 IU hCG is administered 24-48 hours after the last GONAL-f injection to induce final follicular maturation.

Down-regulation with a gonadotropin-releasing hormone (GnRH) agonist or antagonist is now commonly used in order to suppress the endogenous LH surge and to control tonic levels of LH. In a commonly used protocol, GONAL-f is started approximately 2 weeks after the start of agonist treatment, both being continued until adequate follicular development is achieved. For example, following two weeks of treatment with an agonist, 150-225 IU GONAL-f are administered for the first 7 days. The dose is then adjusted according to the ovarian response.

Overall experience with IVF indicates that in general the treatment success rate remains stable during the first four attempts and gradually declines thereafter.

Women with anovulation resulting from severe LH and FSH deficiency.

In LH and FSH deficient women (hypogonadotrophic hypogonadism), the objective of GONAL-f therapy in association with lutropin alfa is to develop a single mature Graafian follicle from which the oocyte will be liberated after the administration of human chorionic gonadotropin (hCG). GONAL-f should be given as a course of daily injections simultaneously with lutropin alfa. Since these patients are amenorrhoeic and have low endogenous oestrogen secretion, treatment can commence at any time.

A recommended regimen commences at 75 IU of lutropin alfa daily with 75-150 IU FSH. Treatment should be tailored to the individual patient's response as assessed by measuring follicle size by ultrasound and oestrogen response.

If an FSH dose increase is deemed appropriate, dose adaptation should preferably be after 7-14 day intervals and preferably by 37.5-75 IU increments. It may be acceptable to extend the duration of stimulation in any one cycle to up to 5 weeks.

When an optimal response is obtained, a single injection of 250 micrograms r-hCG or 5,000 IU up to 10,000 IU hCG should be administered 24-48 hours after the last GONAL-f and lutropin alfa injections. The patient is recommended to have coitus on the day of, and on the day following, hCG administration.

Alternatively, IUI may be performed.

Luteal phase support may be considered since lack of substances with luteotrophic activity (LH/hCG) after ovulation may lead to premature failure of the corpus luteum.

If an excessive response is obtained, treatment should be stopped and hCG withheld. Treatment should recommence in the next cycle at a dose of FSH lower than that of the previous cycle.

Men with hypogonadotrophic hypogonadism

GONAL-f should be given at a dose of 150 IU three times a week, concomitantly with hCG, for a minimum of 4 months. If after this period, the patient has not responded, the combination treatment may be continued; current clinical experience indicates that treatment for at least 18 months may be necessary to achieve spermatogenesis.

Special population

Older people

There is no relevant use of GONAL-f in the elderly population. Safety and effectiveness of GONAL-f in elderly patients have not been established.

Renal or hepatic impairment

Safety, efficacy and pharmacokinetics of GONAL-f in patients with renal or hepatic impairment have not been established.

Paediatric population

There is no relevant use of GONAL-f in the paediatric population.

Method of administration

GONAL-f is intended for subcutaneous use. The first injection of GONAL-f should be performed under direct medical supervision. Self-administration of GONAL-f should only be performed by patients who are well motivated, adequately trained and have access to expert advice.

As GONAL-f pre-filled pen with multidose cartridge is intended for several injections, clear instructions should be provided to the patients to avoid misuse of the multidose presentation.

For instructions on the administration with the pre-filled pen, see section 6.6 and the "Instructions for Use".

4.3 Contraindications

- hypersensitivity to the active substance or to any of the excipients listed in section 6.1
- tumours of the hypothalamus or pituitary gland
- ovarian enlargement or ovarian cyst not due to polycystic ovarian syndrome
- gynaecological haemorrhages of unknown aetiology
- ovarian, uterine or mammary carcinoma

GONAL-f must not be used when an effective response cannot be obtained, such as:

- primary ovarian failure
- malformations of sexual organs incompatible with pregnancy
- fibroid tumours of the uterus incompatible with pregnancy
- primary testicular insufficiency

4.4 Special warnings and precautions for use

GONAL-f is a potent gonadotrophic substance capable of causing mild to severe adverse reactions, and should only be used by physicians who are thoroughly familiar with infertility problems and their management.

Gonadotropin therapy requires a certain time commitment by physicians and supportive health professionals, as well as the availability of appropriate monitoring facilities. In women, safe and effective use of GONAL-f calls for monitoring of ovarian response with ultrasound, alone or preferably in combination with measurement of serum oestradiol levels, on a regular basis. There may be a degree of inter-patient variability in response to FSH administration, with a poor response to FSH in some patients and exaggerated response in others. The lowest effective dose in relation to the treatment objective should be used in both men and women.

Porphyria

Patients with porphyria or a family history of porphyria should be closely monitored during treatment with GONAL-f. Deterioration or a first appearance of this condition may require cessation of treatment.

Treatment in women

Before starting treatment, the couple's infertility should be assessed as appropriate and putative contraindications for pregnancy evaluated. In particular, patients should be evaluated for hypothyroidism, adrenocortical deficiency, hyperprolactinemia and appropriate specific treatment given.

Patients undergoing stimulation of follicular growth, whether as treatment for anovulatory infertility or ART procedures, may experience ovarian enlargement or develop hyperstimulation. Adherence to recommended GONAL-f dose and regimen of administration, and careful monitoring of therapy will minimise the incidence of such events. For accurate interpretation of the indices of follicle development and maturation, the physician should be experienced in the interpretation of the relevant tests.

In clinical trials, an increase of the ovarian sensitivity to GONAL-f was shown when administered with lutropin alfa. If an FSH dose increase is deemed appropriate, dose adaptation should preferably be at 7-14 day intervals and preferably with 37.5-75 IU increments.

No direct comparison of GONAL-f/LH versus human menopausal gonadotropin (hMG) has been performed. Comparison with historical data suggests that the ovulation rate obtained with GONAL-f/LH is similar to that obtained with hMG.

Ovarian Hyperstimulation Syndrome (OHSS)

A certain degree of ovarian enlargement is an expected effect of controlled ovarian stimulation. It is more commonly seen in women with polycystic ovarian syndrome and usually regresses without treatment.

In distinction to uncomplicated ovarian enlargement, OHSS is a condition that can manifest itself with increasing degrees of severity. It comprises marked ovarian enlargement, high serum sex steroids, and an increase in vascular permeability which can result in an accumulation of fluid in the peritoneal, pleural and, rarely, in the pericardial cavities.

The following symptomatology may be observed in severe cases of OHSS: abdominal pain, abdominal distension, severe ovarian enlargement, weight gain, dyspnoea, oliguria and gastrointestinal symptoms including nausea, vomiting and diarrhoea. Clinical evaluation may reveal hypovolaemia, haemoconcentration, electrolyte imbalances, ascites, haemoperitoneum, pleural effusions, hydrothorax, or acute pulmonary distress. Very rarely, severe OHSS may be complicated by ovarian torsion or thromboembolic events such as pulmonary embolism, ischaemic stroke or myocardial infarction.

Independent risk factors for developing OHSS include polycystic ovarian syndrome high absolute or rapidly rising serum oestradiol levels (e.g. > 900 pg/mL or > 3,300 pmol/L in anovulation; > 3,000 pg/mL or > 11,000 pmol/L in ART) and large number of developing ovarian follicles (e.g. > 3 follicles of \geq 14 mm in diameter in anovulation; \geq 20 follicles of \geq 12 mm in diameter in ART).

Adherence to recommended GONAL-f dose and regimen of administration can minimise the risk of ovarian hyperstimulation (see sections 4.2 and 4.8). Monitoring of stimulation cycles by ultrasound scans as well as oestradiol measurements are recommended to early identify risk factors.

There is evidence to suggest that hCG plays a key role in triggering OHSS and that the syndrome may be more severe and more protracted if pregnancy occurs. Therefore, if signs of ovarian hyperstimulation occur such as serum oestradiol level > 5,500 pg/mL or > 20,200 pmol/L and/or \geq 40 follicles in total, it is recommended that hCG be withheld and the patient be advised to refrain from coitus or to use barrier contraceptive methods for at least 4 days. OHSS may progress rapidly (within 24 hours) or over several days to become a serious medical event. It most often occurs after hormonal treatment has been discontinued and reaches its maximum at about seven to ten days following treatment. Therefore patients should be followed for at least two weeks after hCG administration.

In ART, aspiration of all follicles prior to ovulation may reduce the occurrence of hyperstimulation.

Mild or moderate OHSS usually resolves spontaneously. If severe OHSS occurs, it is recommended that gonadotropin treatment be stopped if still ongoing, and that the patient be hospitalised and appropriate therapy be started.

Multiple pregnancy

In patients undergoing ovulation induction, the incidence of multiple pregnancy is increased compared with natural conception. The majority of multiple conceptions are twins. Multiple pregnancy, especially of high order, carries an increased risk of adverse maternal and perinatal outcomes.

To minimise the risk of multiple pregnancy, careful monitoring of ovarian response is recommended.

In patients undergoing ART procedures the risk of multiple pregnancy is related mainly to the number of embryos replaced, their quality and the patient age.

The patients should be advised of the potential risk of multiple births before starting treatment.

Pregnancy loss

The incidence of pregnancy loss by miscarriage or abortion is higher in patients undergoing stimulation of follicular growth for ovulation induction or ART than following natural conception.

Ectopic pregnancy

Women with a history of tubal disease are at risk of ectopic pregnancy, whether the pregnancy is obtained by spontaneous conception or with fertility treatments. The prevalence of ectopic pregnancy after ART, was reported to be higher than in the general population.

Reproductive system neoplasms

There have been reports of ovarian and other reproductive system neoplasms, both benign and malignant, in women who have undergone multiple treatment regimens for infertility treatment. It is not yet established whether or not treatment with gonadotropins increases the risk of these tumours in infertile women.

Congenital malformation

The prevalence of congenital malformations after ART may be slightly higher than after spontaneous conceptions. This is thought to be due to differences in parental characteristics (e.g. maternal age, sperm characteristics) and multiple pregnancies.

Thromboembolic events

In women with recent or ongoing thromboembolic disease or women with generally recognised risk factors for thromboembolic events, such as personal or family history, treatment with gonadotropins may further increase the risk for aggravation or occurrence of such events. In these women, the benefits of gonadotropin administration need to be weighed against the risks. It should be noted however that pregnancy itself as well as OHSS also carry an increased risk of thromboembolic events.

Treatment in men

Elevated endogenous FSH levels are indicative of primary testicular failure. Such patients are unresponsive to GONAL-f/hCG therapy. GONAL-f should not be used when an effective response cannot be obtained.

Semen analysis is recommended 4 to 6 months after the beginning of treatment as part of the assessment of the response.

Sodium content

GONAL-f contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially “sodium-free”.

4.5 Interaction with other medicinal products and other forms of interaction

Concomitant use of GONAL-f with other medicinal products used to stimulate ovulation (e.g. hCG, clomiphene citrate) may potentiate the follicular response, whereas concurrent use of a GnRH agonist or antagonist to induce pituitary desensitisation may increase the dose of GONAL-f needed to elicit an adequate ovarian response. No other clinically significant medicinal product interaction has been reported during GONAL-f therapy.

4.6 Fertility, pregnancy and lactation

Pregnancy

There is no indication for use of GONAL-f during pregnancy. Data on a limited number of exposed pregnancies (less than 300 pregnancy outcomes) indicate no malformative or feto/neonatal toxicity of follitropin alfa.

No teratogenic effect has been observed in animal studies (see section 5.3).

In case of exposure during pregnancy, clinical data are not sufficient to exclude a teratogenic effect of GONAL-f.

Breast-feeding

GONAL-f is not indicated during breast-feeding.

Fertility

GONAL-f is indicated for use in infertility (see section 4.1).

4.7 Effects on ability to drive and use machines

GONAL-f is expected to have no or negligible influence on the ability to drive and use machines.

4.8 Undesirable effects

Summary of the safety profile

The most commonly reported adverse reactions are headache, ovarian cysts and local injection site reactions (e.g. pain, erythema, haematoma, swelling and/or irritation at the site of injection).

Mild or moderate ovarian hyperstimulation syndrome (OHSS) has been commonly reported and should be considered as an intrinsic risk of the stimulation procedure. Severe OHSS is uncommon (see section 4.4).

Thromboembolism may occur very rarely (see section 4.4).

List of adverse reactions

The following definitions apply to the frequency terminology used hereafter: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), very rare ($< 1/10,000$).

Treatment in women

Immune system disorders

Very rare: Mild to severe hypersensitivity reactions including anaphylactic reactions and shock

Nervous system disorders

Very common: Headache

Vascular disorders

Very rare: Thromboembolism (both in association with and separate from OHSS)

Respiratory, thoracic and mediastinal disorders

Very rare: Exacerbation or aggravation of asthma

Gastrointestinal disorders

Common: Abdominal pain, abdominal distension, abdominal discomfort, nausea, vomiting, diarrhoea

Reproductive system and breast disorders

Very common: Ovarian cysts

Common: Mild or moderate OHSS (including associated symptomatology)

Uncommon: Severe OHSS (including associated symptomatology) (see section 4.4)

Rare: Complication of severe OHSS

General disorders and administration site conditions

Very common: Injection site reactions (e.g. pain, erythema, haematoma, swelling and/or irritation at the site of injection)

Treatment in men

Immune system disorders

Very rare: Mild to severe hypersensitivity reactions including anaphylactic reactions and shock

Respiratory, thoracic and mediastinal disorders

Very rare: Exacerbation or aggravation of asthma

Skin and subcutaneous tissue disorders

Common: Acne

Reproductive system and breast disorders

Common: Gynaecomastia, Varicocele

General disorders and administration site conditions

Very common: Injection site reactions (e.g. pain, erythema, haematoma, swelling and/or irritation at the site of injection)

Investigations

Common: Weight gain

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

4.9 Overdose

The effects of an overdose of GONAL-f are unknown, nevertheless, there is a possibility that OHSS may occur (see section 4.4).

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Sex hormones and modulators of the genital systems, gonadotropins, ATC code: G03GA05.

In women, the most important effect resulting from parenteral administration of FSH is the development of mature Graafian follicles. In women with anovulation, the object of GONAL-f therapy is to develop a single mature Graafian follicle from which the ovum will be liberated after the administration of hCG.

Clinical efficacy and safety in women

In clinical trials, patients with severe FSH and LH deficiency were defined by an endogenous serum LH level < 1.2 IU/L as measured in a central laboratory. However, it should be taken into account that there are variations between LH measurements performed in different laboratories.

In clinical studies comparing r-hFSH (follitropin alfa) and urinary FSH in ART (see table below) and in ovulation induction, GONAL-f was more potent than urinary FSH in terms of a lower total dose and a shorter treatment period needed to trigger follicular maturation.

In ART, GONAL-f at a lower total dose and shorter treatment period than urinary FSH, resulted in a higher number of oocytes retrieved when compared to urinary FSH.

Table: Results of study GF 8407 (randomised parallel group study comparing efficacy and safety of GONAL-f with urinary FSH in assisted reproduction technologies)

	GONAL-f (n = 130)	urinary FSH (n = 116)
Number of oocytes retrieved	11.0 ± 5.9	8.8 ± 4.8
Days of FSH stimulation required	11.7 ± 1.9	14.5 ± 3.3
Total dose of FSH required (number of FSH 75 IU ampoules)	27.6 ± 10.2	40.7 ± 13.6
Need to increase the dose (%)	56.2	85.3

Differences between the 2 groups were statistically significant ($p < 0.05$) for all criteria listed.

Clinical efficacy and safety in men

In men deficient in FSH, GONAL-f administered concomitantly with hCG for at least 4 months induces spermatogenesis.

5.2 Pharmacokinetic properties

Following intravenous administration, follitropin alfa is distributed to the extracellular fluid space with an initial half-life of around 2 hours and eliminated from the body with a terminal half-life of about one day. The steady state volume of distribution and total clearance are 10 l and 0.6 l/h, respectively. One-eighth of the follitropin alfa dose is excreted in the urine.

Following subcutaneous administration, the absolute bioavailability is about 70 %. Following repeated administration, follitropin alfa accumulates 3-fold achieving a steady-state within 3-4 days. In women whose endogenous gonadotropin secretion is suppressed, follitropin alfa has nevertheless been shown to effectively stimulate follicular development and steroidogenesis, despite unmeasurable LH levels.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of single and repeated dose toxicity and genotoxicity additional to that already stated in other sections of this SmPC.

Impaired fertility has been reported in rats exposed to pharmacological doses of follitropin alfa (≥ 40 IU/kg/day) for extended periods, through reduced fecundity.

Given in high doses (≥ 5 IU/kg/day) follitropin alfa caused a decrease in the number of viable foetuses without being a teratogen, and dystocia similar to that observed with urinary Menopausal Gonadotropin (hMG). However, since GONAL-f is not indicated in pregnancy, these data are of limited clinical relevance.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Poloxamer 188
Sucrose
Methionine
Sodium dihydrogen phosphate monohydrate
Disodium phosphate dihydrate
m-Cresol
Phosphoric acid, concentrated
Sodium hydroxide
Water for injections

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

2 years.

Once opened, the medicinal product may be stored for a maximum of 28 days at or below 25°C. The patient should write on the GONAL-f pre-filled pen the day of the first use.

6.4 Special precautions for storage

Store in a refrigerator (2°C-8°C). Do not freeze.

Before opening and within its shelf life, the medicinal product may be removed from the refrigerator, without being refrigerated again, for up to 3 months at or below 25°C. The product must be discarded if it has not been used after 3 months.

Store in the original package, in order to protect from light.

For in-use storage conditions, see section 6.3.

6.5 Nature and contents of container

0.75 mL of solution for injection in 3 mL cartridge (Type I glass), with a plunger stopper (halobutyl rubber) and an aluminium crimp cap with a black rubber inlay.

Pack of one pre-filled pen and 12 needles to be used with the pen for administration.

6.6 Special precautions for disposal and other handling

See the “Instructions for Use”.

The solution should not be administered if it contains particles or is not clear.

Any unused solution must be discarded not later than 28 days after first opening.

GONAL-f 450 IU/0.75 mL (33 micrograms/0.75 mL) is not designed to allow the cartridge to be removed.

Discard used needles immediately after injection.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Merck Europe B.V.

Gustav Mahlerplein 102

1082 MA Amsterdam

The Netherlands

Package leaflet: Information for the user

GONAL-f 450 IU/0.75 mL (33 micrograms/0.75 mL) solution for injection in a pre-filled pen Follitropin alfa

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What GONAL-f is and what it is used for
2. What you need to know before you use GONAL-f
3. How to use GONAL-f
4. Possible side effects
5. How to store GONAL-f
6. Contents of the pack and other information
Instructions for Use.

1. What GONAL-f is and what it is used for

What GONAL-f is

GONAL-f contains a medicine called “follitropin alfa”. Follitropin alfa is a type of “Follicle Stimulating Hormone” (FSH) which belongs to the family of hormones called “gonadotropins”. Gonadotropins are involved in reproduction and fertility.

What GONAL-f is used for

In adult women, GONAL-f is used:

- to help release an egg from the ovary (ovulation) in women that cannot ovulate and that did not respond to treatment with a medicine called “clomiphene citrate”.
- together with another medicine called “lutropin alfa” (“Luteinising Hormone” or LH) to help release egg from the ovary (ovulation) in women that are not ovulating because their body is producing very little gonadotropins (FSH and LH).
- to help develop several follicles (each containing an egg) in women undergoing assisted reproductive technology procedures (procedures that may help you to become pregnant) such as “*in vitro* fertilisation”, “gamete intra-fallopian transfer” or “zygote intra-fallopian transfer”.

In adult men, GONAL-f is used:

- together with another medicine called “human Chorionic Gonadotropin” (hCG) to help produce sperm in men that are infertile due to a low level of certain hormones.

2. What you need to know before you use GONAL-f

You and your partner’s fertility should be evaluated before the treatment is started by a doctor experienced in treating fertility disorders.

Do not use GONAL-f

- if you are allergic to Follicle Stimulating Hormone or any of the other ingredients of this medicine listed in section 6.
- if you have a tumour in your hypothalamus or pituitary gland (both are parts of the brain).
- If you are **a woman**:
 - with large ovaries or sacs of fluids within the ovaries (ovarian cysts) of unknown origin.
 - with unexplained vaginal bleeding.
 - with cancer in your ovaries, womb or breasts.
 - with a condition that usually makes normal pregnancy impossible, such as ovarian failure (early menopause), or malformed reproductive organs.
- If you are **a man**:
 - with damaged testicles that cannot be healed.

Do not use GONAL-f if any of the above applies to you. If you are not sure, talk to your doctor before using this medicine.

Warnings and precautions

Porphyria

Tell your doctor before you start treatment, if you or any member of your family have porphyria (an inability to break down porphyrins that may be passed on from parents to children).

Tell your doctor straight away if:

- your skin becomes fragile and easily blistered, especially skin that has been frequently in the sun, and/or
- you have stomach, arm or leg pain.

In case of the above events your doctor may recommend that you stop treatment.

Ovarian Hyper-Stimulation Syndrome (OHSS)

If you are a woman, this medicine increases your risk of developing OHSS. This is when your follicles develop too much and become large cysts. If you get lower abdominal pain, gain any weight rapidly, feel sick or are vomiting or if you have difficulty in breathing, talk to your doctor straight away who might ask you to stop using this medicine (see section 4).

In case you are not ovulating, and if the recommended dose and schedule of administration are adhered to, the occurrence of OHSS is less likely. GONAL-f treatment seldom causes severe OHSS unless the medicine that is used for final follicular maturation (containing human Chorionic Gonadotropin, hCG) is administered. If you are developing OHSS your doctor may not give you any hCG in this treatment cycle and you may be told not to have sex or to use a barrier contraceptive method for at least four days.

Multiple pregnancy

When using GONAL-f, you have a higher risk of being pregnant with more than one child at the same time (“multiple pregnancy”, mostly twins), than if you conceived naturally. Multiple pregnancy may lead to medical complications for you and your babies. You can reduce the risk of multiple pregnancy by using the right dose of GONAL-f at the right times. When undergoing assisted reproductive technology the risk of having a multiple pregnancy is related to your age, the quality and the number of fertilised eggs or embryos placed inside you.

Miscarriage

When undergoing assisted reproductive technology or stimulation of your ovaries to produce eggs, you are more likely to have a miscarriage than the average woman.

Blood clotting problems (thromboembolic events)

If you had in the past or recently blood clots in the leg or in the lung, or a heart attack or stroke, or if those happened in your family, then you might have a higher risk that these problems occur or become worse with GONAL-f treatment.

Men with too much FSH in their blood

If you are a man, having too much FSH in your blood can be a sign of damaged testicles. GONAL-f usually does not work if you have this problem.

If your doctor decides to try GONAL-f treatment, to monitor the treatment, they may ask you to provide semen for analysis 4 to 6 months after starting treatment.

Children

GONAL-f is not indicated for use in children.

Other medicines and GONAL-f

Tell your doctor if you are taking, have recently taken or might take any other medicines.

- If you use GONAL-f with other medicines which help ovulation (such as hCG or clomiphene citrate), this may increase the response of your follicles.
- If you use GONAL-f at the same time as a “gonadotropin-releasing hormone” (GnRH) agonist or antagonist (these medicines reduce your sex hormone levels and stop you ovulating) you may need a higher dose of GONAL-f to produce follicles.

Pregnancy and breast-feeding

Do not use GONAL-f if you are pregnant or breast-feeding.

Driving and using machines

It is not expected that this medicine will affect your ability to drive and use machines.

GONAL-f contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per dose, therefore it is considered essentially “sodium-free”.

3. How to use GONAL-f

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Using this medicine

- GONAL-f is intended to be given by injection just under the skin (subcutaneously). The pre-filled pen can be used for several injections.

- The first injection of GONAL-f should be given under supervision of your doctor.
- Your doctor or nurse will show you how to use the GONAL-f pre-filled pen to inject the medicine.
- If you administer GONAL-f to yourself, please carefully read and follow the “Instructions for Use”.

How much to use

Your doctor will decide how much medicine you will take and how often. The doses described below are stated in International Units (IU) and millilitres (mL).

Women

If you are not ovulating and have irregular or no periods.

- GONAL-f is usually given every day.
- If you have irregular periods, start using GONAL-f within the first 7 days of your menstrual cycle. If you do not have periods you can start using the medicine on any convenient day.
- The usual starting dose of GONAL-f is 75 to 150 IU (0.12 to 0.24 mL) each day.
- Your dose of GONAL-f may be increased every 7 or every 14 days by 37.5 to 75 IU, until you get the desired response.
- The maximum daily dose of GONAL-f is usually not higher than 225 IU (0.36 mL).
- When you get the desired response, you will be given a single injection of 250 micrograms of “recombinant hCG” (r-hCG, an hCG made in a laboratory by a special DNA technique), or 5,000 to 10,000 IU of hCG, 24 to 48 hours after your last GONAL-f injection. The best time to have sex is on the day of the hCG injection and the day after.

If your doctor cannot see a desired response after 4 weeks, that treatment cycle with GONAL-f should be stopped. For the following treatment cycle, your doctor will give you a higher starting dose of GONAL-f than before.

If your body responds too strongly, your treatment will be stopped and you will not be given any hCG (see section 2, OHSS). For the following cycle, your doctor will give you a lower dose of GONAL-f than before.

If you are not ovulating, having no periods and have been diagnosed with very low levels of FSH and LH hormones

- The usual starting dose of GONAL-f is 75 to 150 IU (0.12 to 0.24 mL) together with 75 IU (0.12 mL) of lutropin alfa.
- You will use these two medicines each day for up to five weeks.
- Your dose of GONAL-f may be increased every 7 or every 14 days by 37.5 to 75 IU, until you get the desired response.
- When you get the desired response, you will be given a single injection of 250 micrograms of “recombinant hCG” (r-hCG, an hCG made in a laboratory by a special DNA technique), or 5,000 to 10,000 IU of hCG, 24 to 48 hours after your last injections of GONAL-f and lutropin alfa. The best time to have sex is on the day of the hCG injection and the day after. Alternatively, intrauterine insemination may be performed by placing the sperm into the womb cavity.

If your doctor cannot see a response after 5 weeks, that treatment cycle with GONAL-f should be stopped. For the following cycle, your doctor will give you a higher starting dose of GONAL-f than before.

If your body responds too strongly, your treatment with GONAL-f will be stopped and you will not be given any hCG (see section 2, OHSS). For the following cycle, your doctor will give you a lower dose of GONAL-f than before.

If you need to develop several eggs for collection prior to any assisted reproductive technology

- The usual starting dose of GONAL-f is 150 to 225 IU (0.24 to 0.36 mL) each day, from day 2 or 3 of your treatment cycle.
- GONAL-f dose may be increased, depending on your response. The maximum daily dose is 450 IU (0.72 mL).
- Treatment is continued until your eggs have developed to a desired point. This usually takes about 10 days but can take any time between 5 and 20 days. Your doctor will use blood tests and/or an ultrasound machine to check when this is.
- When your eggs are ready, you will be given a single injection of 250 micrograms “recombinant hCG” (r-hCG, an hCG made in a laboratory by a special recombinant DNA technique), or 5,000 IU to 10,000 IU of hCG, 24 to 48 hours after the last GONAL-f injection. This gets your eggs ready for collection.

In other cases, your doctor may first stop you from ovulating by using a gonadotropin-releasing hormone (GnRH) agonist or antagonist. Then GONAL-f is started approximately two weeks after start of agonist treatment. The GONAL-f and GnRH agonist are then both given until your follicles develop as desired. For example, after two weeks of GnRH agonist treatment, 150 to 225 IU GONAL-f is administered for 7 days. The dose is then adjusted according to your ovarian response.

Men

- The usual dose of GONAL-f is 150 IU (0.24 mL) together with hCG.
- You will use these two medicines three times a week for at least 4 months.
- If you have not responded to treatment after 4 months, your doctor may suggest that you continue using these two medicines for at least 18 months.

If you use more GONAL-f than you should

The effects of taking too much GONAL-f are unknown. Nevertheless one could expect Ovarian Hyper-Stimulation Syndrome (OHSS) to occur, which is described in section 4. However the OHSS will only occur if hCG is also administered (see section 2, OHSS).

If you forget to use GONAL-f

If you forget to use GONAL-f, do not take a double dose to make up for a forgotten dose. Please talk to your doctor as soon as you notice that you forgot a dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Serious side effects in women

- Lower abdominal pain together with nausea or vomiting may be the symptoms of Ovarian Hyper-Stimulation Syndrome (OHSS). This may indicate that the ovaries over-reacted to the treatment and that large ovarian cysts developed (see also in section 2. under “Ovarian Hyper-Stimulation Syndrome”). This side effect is common (may affect up to 1 in 10 people).

- The OHSS may become severe with clearly enlarged ovaries, decreased urine production, weight gain, difficulty in breathing and/or possible fluid accumulation in your stomach or chest. This side effect is uncommon (may affect up to 1 in 100 people).
- Complications of OHSS such as twisting of ovaries or blood clotting may occur rarely (may affect up to 1 in 1,000 people).
- Serious blood clotting complications (thromboembolic events) sometimes independent of OHSS may be found very rarely (may affect up to 1 in 10,000 people). This could cause chest pain, breathlessness, stroke or heart attack (see also in section 2. under “Blood clotting problems”).

Serious side effects in men and women

- Allergic reactions such as rash, red skin, hives, swelling of your face with difficulty breathing can sometimes be serious. This side effect is very rare (may affect up to 1 in 10,000 people).

If you notice any of the above-listed side effects you should immediately contact your doctor who might ask you to stop using GONAL-f.

Other side effects in women

Very common (may affect more than 1 in 10 people):

- Sacs of fluid within the ovaries (ovarian cysts)
- Headache
- Local reactions at the injection site, such as pain, redness, bruising, swelling and/or irritation

Common (may affect up to 1 in 10 people):

- Abdominal pain
- Feeling sick, vomiting, diarrhoea, abdominal cramps and bloating

Very rare (may affect up to 1 in 10,000 people):

- Allergic reactions such as rash, red skin, hives, swelling of your face with difficulty breathing may occur. These reactions can sometimes be serious.
- Your asthma may get worse

Other side effects in men

Very common (may affect more than 1 in 10 people):

- Local reactions at the injection site, such as pain, redness, bruising, swelling and/or irritation

Common (may affect up to 1 in 10 people):

- Swelling of the veins above and behind the testicles (a varicocele).
- Breast development, acne or weight gain.

Very rare (may affect up to 1 in 10,000 people):

- Allergic reactions such as rash, red skin, hives, swelling of your face with difficulty in breathing may occur. These reactions can sometimes be serious.
- Your asthma may get worse.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store GONAL-f

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the cartridge label or carton after EXP. The expiry date refers to the last day of that month.

Store in a refrigerator (2°C - 8°C). Do not freeze.

Within its shelf life, the product may be stored at or below 25°C for up to 3 months without being refrigerated again and must be discarded if it has not been used after 3 months.

Store in the original package in order to protect from light.

Do not use GONAL-f if you notice any visible signs of deterioration, if the liquid contains particles or is not clear.

Please write on the GONAL-f pre-filled pen the day you first use it.

- Once opened, it may be stored for a maximum of 28 days outside of the fridge (at or below 25°C).
- Do not use any medicine left in your pre-filled pen after 28 days.

At the end of the treatment any unused solution must be discarded.

Do not throw away any medicines via wastewater. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What GONAL-f contains

- The active substance is follitropin alfa.
- There are 600 IU (44 micrograms) of follitropin alfa in each millilitre of liquid. Each pre-filled pen with multidose cartridge delivers 450 IU (33 micrograms) in 0.75 mL.
- The other ingredients are poloxamer 188, sucrose, methionine, sodium dihydrogen phosphate monohydrate, disodium phosphate dihydrate, m-cresol, concentrated phosphoric acid, sodium hydroxide and water for injections.

What GONAL-f looks like and contents of the pack

- GONAL-f is presented as a clear, colourless liquid for injection in a pre-filled pen.
- It is supplied in packs with 1 pre-filled pen and 12 disposable needles.

Marketing Authorisation Holder

Merck Europe B.V., Gustav Mahlerplein 102, 1082 MA Amsterdam, The Netherlands.

GONAL-f Pre-filled pen 450 IU/0.75 mL

Instructions for Use

Contents

1. How to use the GONAL-f pre-filled pen
2. How to use your GONAL-f pre-filled pen Treatment diary
3. Before you start using your GONAL-f pre-filled pen
4. Getting your GONAL-f pre-filled pen ready for injection
5. Setting the dose prescribed by your doctor
6. Injecting the dose
7. After the injection
8. GONAL-f pre-filled pen Treatment diary (see table at the end)

Warning: Please read these instructions for use before using your GONAL-f pre-filled pen. Follow the procedure exactly, as it may differ from your past experience.

1. How to use the GONAL-f pre-filled pen

- Do not share the pen. The pen is for subcutaneous injection only.
- The numbers on the **Dose Feedback Window** are measured in International Units or IUs. Your doctor will have told you how many IUs to inject each day.
- The numbers displayed in the **Dose Feedback Window** help you to:
 - a. Dial your prescribed dose.



- b. Verify a complete injection.



- c. Read the dose remaining to be injected with a second pen.



- Give yourself the injection at the same time each day. Example: 
- Your doctor/pharmacist will tell you how many pens you need to complete your treatment.

2. How to use your GONAL-f pre-filled pen Treatment Diary

A treatment diary is included on the last page.

Use the treatment diary to record the amount of injected IUs you use each time.

- Record the treatment day number (1), date (2) and time (3) of your injection.
- In the first line of the table the volume of your pen is already recorded for you (4).
- Record your prescribed dose in the ‘‘Prescribed Dose’’ section (5).
- Verify you dialed the right dose before injecting (6).
- After injection, read the number shown in the **Dose Feedback Window**.

- Confirm you receive a complete injection (7) or record the number shown in the **Dose Feedback Window** if other than ‘0’ (8).
- When needed, inject yourself using a second pen, dialing your remaining dose written in the ‘Amount Displayed After Injection’ section (8).
- Record this remaining dose in the ‘Amount Set to Inject’ section in the next row (6).

CAUTION:
Using your treatment diary to record your daily injection(s) allows you to verify every day that you received the full prescribed dose.


An example of a treatment diary:

1 Treatment Day Number	2 Date	3 Time	4 Pen Volume 450 IU/0.75 mL	5 Prescribed Dose	6 7 8		
					Dose Feedback Window		
					Amount Set to Inject	Amount Displayed After Injection	
#1	10/06	07:00	450 IU	175	175	<input checked="" type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen
#2	11/06	07:00	450 IU	175	175	<input checked="" type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen
#3	12/06/	07:00	450 IU	175	175	<input type="checkbox"/> if "0", injection complete	<input checked="" type="checkbox"/> if not "0", need second injection Inject this amount .75 using new pen
#3	12/06	07:00	450 IU	N/A	75	<input checked="" type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen

Note: The 300 IU pen maximum single-dose setting is 300 IU; the 450 IU pen maximum single-dose setting is 450 IU; the 900 IU pen maximum single-dose setting is 450 IU.

3. Before you start using your GONAL-f pre-filled pen

- Wash your hands with soap and water.
- Find a clean area and a **flat surface**.
- Verify the **expiration date** on the pen label.
- Gather everything you need and lay it out:




- | | | |
|--------------------------------|------------------------------|-------------------------------|
| 1. Dose setting knob | 5. Threaded needle connector | 9. Inner needle shield |
| 2. Dose Feedback Window | 6. Pen cap | 10. Outer needle cap |
| 3. Plunger piston | 7. Peel-off seal tab | 11. Alcohol swabs |
| 4. Reservoir holder | 8. Removable needle | 12. Sharps disposal container |

4. Getting your GONAL-f pre-filled pen ready for injection

4.1. Take off the pen cap

4.2. Verify that the Dose Feedback Window is set to "0".



4.3. Prepare your needle for Injection

- Get a new needle - only use the "single-use" needles supplied.
- Hold the outer needle cap firmly.
- Check that the peel-off seal on the outer needle cap is not damaged or loose:

Example of a good seal



Example of a bad seal



- Remove the peel-off seal.



CAUTION:


If the peel-off seal is damaged or loose, do not use the needle. Throw it away in a sharps disposal container. Get a new needle.

4.4. Attach the needle

- Screw the threaded tip of the GONAL-f pre-filled pen into the outer needle cap until you feel a light resistance.
Warning: Do not attach the needle too tightly; the needle could be difficult to remove after the injection.
- Remove the outer needle cap by pulling it gently.



Put it aside for later use.



- Hold the GONAL-f pre-filled pen with the needle pointing upward.
- Carefully remove and discard the green inner shield.




4.5. Look closely at the tip of the needle for tiny drop(s) of fluid

- If you see a tiny drop(s) of fluid proceed to **Section 5: Setting the dose prescribed by your doctor.**

Warning:


ONLY check for drop(s) the **FIRST TIME** you use a new GONAL-f pre-filled pen to remove air in the system.



CAUTION:
*If you do not see a tiny drop(s) at or near the needle tip **the first time** you use a new pen, you must perform the steps on the next page.*


If you do not see a tiny drop(s) of fluid at or near the tip the first time you use a new pen:

1. Gently turn the dose setting knob clockwise until it **reads 25** in the Dose Feedback Window. You can turn the dose knob backwards if you turn it past 25.



2. Hold the pen with the needle pointing upward.
3. Tap the reservoir holder gently.


4. Press the dose setting knob **as far as it will go**. A tiny drop of liquid will appear at the tip of the needle.
5. Verify that the Dose-Feedback Window reads "0".




6. Proceed to **Section 5: Setting the dose prescribed by your doctor**.

5. Setting the dose prescribed by your doctor

- 5.1. The pen contains 450 IU follitropin alpha.
 - The 450 IU pen **maximum single-dose setting is 450 IU** and the smallest single-dose setting is 12.5 IU.
- 5.2. **Turn the dose setting knob until your intended dose shows in the Dose Feedback Window.**



Turn the dose setting knob **forward** to dial up



Turn the dose setting knob **backward** to correct the dose


- 5.3. Set the dose that was prescribed by your doctor (in the example shown in the figure, it is 50 IU).



Warning: Check that the Dose Feedback Window displays your **complete prescribed dose** before you move on to the next step.

6. Injecting the dose

- 6.1. Choose an injection site in the area your doctor or nurse has told you to give the injection.
To minimize skin irritation, select a different injection site each day.




- 6.2. Clean the skin by wiping with an alcohol swab.
- 6.3. Verify once more that the Dose Feedback Window displays the correct dose.
- 6.4. Inject the dose as you were trained to do by your doctor or nurse.


- Slowly push the needle into the skin entirely (1).
- **Press the dose knob down as far as it will go and hold it to complete the full injection.**



- Hold the dose knob down for a minimum of 5 seconds to ensure you inject the full dose (2). The larger the dose, the longer it will take to inject.
- The dose number shown in the Dose Feedback Window will turn back to 0.



- After a minimum of 5 seconds, pull the needle out of the skin while keeping the dose setting knob pressed down (3).
- Release the dose setting knob.



Warning: Always make sure to use a new needle for each injection.

7. After the injection

- 7.1. Verify you have given a complete injection.
 - Check that the Dose Feedback Window shows “0”.



Warning: If the **Dose Feedback Window** shows a number higher than 0, the GONAL-f pre-filled pen is empty and you have not received your full prescribed dose.

7.2. Complete a partial injection (only when needed).



- The **Dose Feedback Window** will indicate the missing amount you need to inject **using a new pen.**



- Repeat Section 3 (“**Before you start using your GONAL-f pre-filled pen** “) through Section 4 (“**Getting your GONAL-f pre-filled pen ready for injection**“) with a second pen.
- Set the dose to the missing amount you recorded in the treatment diary or the number still displayed in the Dose Feedback Window on your previous pen, and inject.

7.3. Removing the needle after each injection.

- Place the outer needle cap on a flat surface.
- Hold the GONAL-f pre-filled pen firmly with one hand, and slip the needle into the outer needle cap.
- Continue by pushing the capped needle against a firm surface until you hear a “click.”
- Grip the outer needle cap and unscrew the needle by **turning anti-clockwise**. Dispose of the used needle safely.
- Never reuse any used needle. Never share needles.
- Recap the pen.



7.4. Storing the GONAL-f pre-filled pen.

CAUTION:

Never store the pen with the needle attached.

Always remove the needle from the GONAL-f pre-filled pen before replacing the pen cap.

- Store the pen in its original packaging in a safe place.
- When the pen is empty, ask your pharmacist how to dispose of it.

Warning: Medicine should not be disposed of via wastewater or household waste.

8. GONAL-f pre-filled pen Treatment Diary

1 Treatment Day Number	2 Date	3 Time	4 Pen Volume 450 IU/0.75 mL	5 Prescribed Dose	6 7 8 Dose Feedback Window		
					Amount Set to Inject	Amount Displayed After Injection	
	/	:	450 IU			<input type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen
	/	:	450 IU			<input type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen
	/	:	450 IU			<input type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen
	/	:	450 IU			<input type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen
	/	:	450 IU			<input type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen -
	/	:	450 IU			<input type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen
	/	:	450 IU			<input type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen
	/	:	450 IU			<input type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen
	/	:	450 IU			<input type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen
	/	:	450 IU			<input type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen
	/	:	450 IU			<input type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen
	/	:	450 IU			<input type="checkbox"/> if "0", injection complete	<input type="checkbox"/> if not "0", need second injection Inject this amountusing new pen

Note: The 300 IU pen maximum single-dose setting is 300 IU; the 450 IU pen maximum single-dose setting is 450 IU; the 900 IU pen maximum single-dose setting is 450 IU.

This instructions for use was last revised in: